

L 82 0 42 84 3000 0 0 (820000000020 0 02 202 10 0



Funded under USAID Cooperative Agreement No. 663-A-00-00-0358-00.

Produced in collaboration with the Ethiopia Public Health Training Initiative, The Carter Center, the Ethiopia Ministry of Health, and the Ethiopia Ministry of Education.

Important Guidelines for Printing and Photocopying

Limited permission is granted free of charge to print or photocopy all pages of this publication for educational, not-for-profit use by health care workers, students or faculty. All copies must retain all author credits and copyright notices included in the original document. Under no circumstances is it permissible to sell or distribute on a commercial basis, or to claim authorship of, copies of material reproduced from this publication.

©2002 by Abraham Alano,

All rights reserved. Except as expressly provided above, no part of this publication may


PREFACE

Nursing is core part in health service delivery system in which health promotion, disease prevention; curative and rehabilitative health strategies are applied. The clinical nursing skills for the nurses are of paramount important not only to provide comprehensive care but also enhance clinical competence. The purpose of preparing this lecture note is to equip nurses with basic clinical nursing skills, which will enable them to dispatch their responsibility as well as to develop uniformity among Ethiopian Professional Nurse Training Higher Institutions.

The lecture note series is designed to have two parts: part-I is composed of most basic clinical skills, where as part two will be covering most advances clinical skills as well as fundamental concepts related to the skills. It is well known that no nursing service can be provided with out basic clinical nursing skills. For nurse to provide health service at different settings; hospital, health center, health post and at the community level including home based care for chronically sick patients, the course is very essential. It is also hoped that other primary and middle level health professional training institution will utilize the lecture notes to rational exercise the professional skills.

The lecture note is therefore organized in logical manner that students can learn from simpler to the complex. It is divided in to units and chapters. Important abbreviations and key terminologies

have been included in order to facilitate teaching learning processes. On top of that learning objectives are clearly stated to indicate the required outcomes. Glossary is prepared at the end to give explanation for terminologies indicated as learning stimulants at beginning of each chapter following the learning objectives. Trial is made to give some scientific explanation for procedure and some relevant study questions are prepared to each chapter to aid students understand of the subject. To enhance systematic approach in conducting nursing care the nursing process is also indicated for most procedures.



ACKNOWLEDGEMENT

My deepest appreciation is to The Carter Center, EPHTI and Professor Dennis Carlson, senior consultant of the Center for his tireless efforts to materialize the issue of staff strengthening and curriculum development. Lecture note preparation is one of the activities that got due attention to strengthen the teaching learning process in Ethiopia by Ethiopian staff. There fore, I congratulate Professor Denis Carlson for the success you achieved with dedicated Ethiopian partners.

I also extend many thanks to colleagues and staff of our School of Nursing particularly Ato Andargachew Kassa and Ato Aweke Yilma for their valuable comments and criticism, other wise the lecture note would have not been shaped this way.

I would like to extend my thanks to Ato Asrat Demissie Academic Vice

Similarly, my felt thanks is to supportive staff of The Carter Center EPHTI, Dr Hailu Yenenh, Ato Aklilu Mulugetta , W/rt Meseret Tsegaw and all other staff for their devoted support through the preparation of this lecture note.

Next, I thank all our College authorities for permission to work on this lecture note besides the routine activities of the college.

Last but not least, my deepest gratitude is to W/rt Lemlem Adebabay who has managed to write the document with in very urgency giving up her rest time and comfort.





TABLE OF CONTENTS

Content	Page
Preface	i
Acknowledgement	iii
Table of Contents.....	v
List of Figures	xii
Abbreviation	xiii
Unit One	
Chapter 1	1
Introduction	1
Definition of nursing	1
Historical background of nursing	2
History of nursing in Ethiopia	8
Nursing process and Critical thinking	10
Unit Two Safety in Health Care facilities	18
Chapter 2: infection control/ universal precaution	18
Nursing process application.....	19
Normal body defense	22
Chain of infection	24
Basic medical asepsis	26

Hand washing	26
Standard precaution	29
Surgical asepsis.....	33
Isolation.....	37
Chapter 3: Care of patient unit.....	49
The patient unit	50
Care of hospital and health care unit equipment.....	54
Unit Three Basic Client Care	58
Chapter 4: Admission, Transfer, and Discharge of client	58
Admission	59
Transfer.....	61
Discharge.....	62
Chapter 5: Vital Signs	66
Definition.....	69
Temperature	69
Pulse	77
Respiration.....	82
Blood pressure.....	83
Chapter 6: Specimen collection	90
General consideration for specimen collection.....	91
Collecting stool specimen	92
Collecting urine specimen.....	94
Collecting sputum	99
Collecting blood specimen.....	101
Chapter 7: Bed making	109
Closed bed	110


Occupied bed.....	110
Post operative bed.....	112
Chapter 8: Personal hygiene and skin care.....	120
Mouth care	120
Bathing.....	125
Bed bath.....	128
Therapeutic bath.....	131
Back care	134
Giving and receiving bedpan and urinals	138
Perineal care.....	140
Hair cares.....	145
Pediculosis treatment	150
Chapter 9: Cold & heat application	158
Care of a patient with fever.....	158
Heat application	159
Cold application	160
Tepid sponge	160

Body positioning	177
Guideline for positioning the client.....	177
Client positioning for examination & treatment.....	179
Crutch walking	183
Application of nursing process.....	183
Teaching technique for crutch walking	184



Enema	236
Cleansing enema	237
Retention enema.....	241
Rectal washout.....	242
Passing flatus tube.....	243
Urinary catheterization	244
Catheterization using straight catheter ...	246
Inserting indwelling catheter.....	249

Uwel



Intravenous therapy	275
Blood transfusion	278
Cut down	283
Administration of vaginal medications..	285
Administration of ophthalmic medication	287
Administration of otic medications	290
Inhalation.....	292
Definition of inhalation.....	292
Oxygen administration	292
Giving oxygen by mask.....	292
Giving oxygen by nasal catheter.....	294
Giving oxygen by tent.....	295
Steam inhalation	297
Nelson's inhalor.....	298
Unit Seven	
Chapter 14 Wound cares	303
Definition	303
Wound healing process.....	304
Dressing a clean wound.....	304
Dressing of septic wound	307
Dressing with a drainage tube.....	309



LIST OF FIGURES


Figure 1. Chain of infection	24
Figure 2. Site for radial pulse measurement	80
Figure 3. Horizontal recumbent position	179
Figure 4. Dorsal recumbent position	179
Figure 5. Prone position	180
Figure 6. Sim position.....	180
Figure 7. Fowler's position	181
Figure 8. Knee-chest position.....	181
Figure 9. Lithotomy position	182
Figure 10. Four –point-gait.....	1825



B.M	Bowel movement
B.M.R.	Basal metabolic rate
B.P	Blood pressure
BPM	Beat perminuli
B.R.P.	Bathroom privilege
BUN	







Q.O.D.	Every other day
R.B.C.	Red blood count or red blood cell
Rh.	Rhesus factor
Rx	Prescription, take
Sol.	Solution
SOS	If necessary
STAT	Immediately -at once
S.C	Subcutaneous
T. I.D	Three times a day
T.P.R.	Temperature, pulse, respiration
Tsp	Teaspoon, tablespoon
U.R.	Upper right
WBC	White blood cells
Wt.	Weight
U.R.Q.	Upper right quadrant
U.L.Q.	Upper lower quadrant
UTI	Urinary tract Infection

UNIT ONE

CHAPTER 1

INTRODUCTION

Learning Objectives:

After completing this unit, the learners will be able to:

- State the modern definition of nursing
- Outline the historical background of nursing world wide and in Ethiopia
- Identify the contribution of significant individuals in nursing
- Describe the nursing process
- Describe critical thinking as an instrument for provision of quality care

Nursing

Definition:

“It is the diagnosis and treatment of human responses to actual or potential health problems” (ANA 1980).

It is assisting the individual, sick or well in the performance of those activities contributing to health or its recovery (to peaceful death) that he will perform unaided, if he had the necessary strength, will or

Nursing is the art and science that involves working with individual, families, and communities to promote wellness of body, mind, and spirit. It is a dynamic, therapeutic and educational process that serves to meet the health needs of the society, including its most vulnerable members.

Historical Background of Nursing

Nursing has a history as long as that of human kind. Human beings have always faced the challenge of fostering health and caring for the ill and dependent. Those who were especially skilled in this area stood out and, in some instances, passed their skills along to others. Uprichard (1973) described the early history of nursing using three images: the folk image, the religious image, and the renaissance image.

The Folk Image of Nursing:

The Nurse as Mother

The early development of nursing was rarely documented, so we must speculate about its character from what we know of early civilizations. The nurse was generally a member of the family or, if not, then a member of the community who demonstrated a special skill in caring for others. Nursing in this perspective was seen largely as a feminine role an extension of mothering. Indeed, the word nursing itself may have been derived from the same root as the words nourish and nurture. This view of nursing was prevalent in the earliest historical records and is still present in primitive cultures.



death; those who worked in them were seen as corrupt and unsavory.

The Emergence of Modern Nursing

To some extent, the three early images of the nurse were held simultaneously for hundreds of years. Then, in the 19th century, one woman changed the course of nursing: **Florence Nightingale**. Although born to wealth and a family well placed in Victorian English Society, Florence Nightingale had a firm belief in Christian ideals that made her disdainful of a life of luxury. She believed her true calling was to minister to the sick. As an intelligent and well-educated woman, she recognized that optimum care of the sick required education. She persevered against family and social opposition and initiated personal study and research into sanitation and health. She studied with **Pastor Fleidner** of 33, was to reorganize the care for the sick at a hospital established for "Gentlewomen in Distressed Circumstances."

Nightingale's success in her first post led Britain's secretary of war to recruit her for a far more arduous reorganization. Britain was then engaged in a major war in the **Crimea**; reports were coming back that more men died of wounds in the hospitals than on the battlefield. Funds were raised and nurses recruited for Florence Nightingale's Crimean campaign. When she arrived at the front, Nightingale found that conditions in the military hospitals were abominable. The absence of sewers and laundry facilities, the lack of supplies, the poor food, and the disorganized medical services

contributed to a death rate of more than 50% among the wounded. Nightingale insisted on retaining control of all of her supplies, funds, and personnel. Her efforts and those of her staff reduced the death rate among the wounded to less than 3%. She eventually completely reformed the military's approach to the health care of the British soldier.

In 1860, she created a school of nursing, which was the model for most nursing education in England. The school was organized around three components: 1) a trained matron with undisputed authority over all members of the staff, 2) a planned course of theoretical and practical training, and 3) a home attached to the hospital in which carefully selected students were placed in the care of "sisters" responsible for their moral and spiritual training. (The English term "**sisters**" used for secular nurses reflects nursing's religious history.) Nightingale established educational standards for the students – she concerned herself not just with health care needs but with human needs.

Her school prepared nurses for hospital care (where they were called "**ward sisters**") and for supervisory and teaching positions. Nightingale also set up a program for preparing "**district**" nurses, the **public health/visiting nurses** of England. She wrote that these district nurses needed additional education because they would be working more independently than the hospital staff members.

Nightingale's strong statements about the role of nurses and their need for lifelong education are still quoted widely today. Perhaps

she, more than anyone else, can be credited with establishing nursing as a profession.

In the early ages, much of the practice of medicine was integrated with religious practices. Before the development of modern nursing, women of nomadic tribes performed nursing duties, such as helping the very young, the old, and the sick, care-dwelling mothers practiced the nursing of their time.

As human needs expanded, nursing development broadened; its interest and functions through the social climates created by religious ideologies, economic development, industrial revolutions, wars, crusades, and education. In this way modern nursing was born.

The intellectual revolution of the 18th and 19th centuries led to a scientific revolution. The dynamic change in economic and political

History of Nursing in Ethiopia

Even though Ethiopia is one of the oldest countries in the world, introduction of modern medicine was very late. Health care of communities and families was by Hakim (wogesha or traditional healers).

Around 1866 missionaries came to Eritrea, (one of the former provinces of Ethiopia) and started to provide medical care for very few members of the society. In 1908 Minlik II hospital was established in the capital of Ethiopia. The hospital was equipped and staffed by Russians.

Later hospital building was continued which raised the need to train health auxiliaries and nurses. In 1949 the Ethiopian Red Cross, School of Nursing was established at Hailesellasie I hospital in Addis Ababa. The training was given for three years. In 1954 Hailesellasie Public Health College was established in Gondar to train health officer, community health nurses and sanitarians to address the health problem of most of the rural population. In line with this, the Centralized school of Nursing formerly under Ministry of health and recently under Addis Ababa University Medical Faculty and Nekemit School of nursing are among the senior nurse's training institutions.

During the regimen of 'Dergue', the former bedside and community health nursing training was changed to comprehensive nursing. An additional higher health professional training institution was also established in Jimma(1983) to train health professionals using educational philosophy of community based and team approach.



NURSING PROCESS and CRITICAL THINKING

Nursing Process:

Definition: Nursing Process is a tool or method for organizing and delivering care or a deliberate intellectual activity where by the practice of nursing is approached in an orderly systematic manner. It is a systematic problem solving approach to client care. It is a series of planned steps and actions directed toward meeting the need and solving problems of people and their significant others; it is systematic, scientific problem solving in action (Sorensen and Luckman, 1986)

Purpose of Nursing Process:

1. To identify clients health care needs
2. To establish nursing care plan so as to meet those needs



health problems. In nursing assessment the best sources of



Physical examination

(Analysis of data is included as part of the assessment. For those who wish to emphasize its importance analysis may be identified as a separate step of the nursing process.)

2. Diagnosis – identification of the following two types of patient problems:

a) Nursing diagnosis – actual or potential health problems that can be managed by independent nursing interventions.

Purposes of the Nursing Diagnosis- the nursing diagnosis serves the following purposes:

- Identifies nursing priorities
- Directs nursing interventions to meet the client's high priority needs
- Provides a common language and forms a basis for communication and understanding between nursing professionals and health care team.
- Guides the formulation of expected outcomes for quality assurance requirements of third party payer.
- Provides a basis for evaluation to determine if nursing

The diagnostic statement

The client may present with more than one problem. Therefore, the nursing diagnosis may be made up of multiple *diagnostic statements*. Each diagnostic statement has two or three parts depending on the healthcare facility. The three-part statement consists of the following components:

- Problem
- Etiology
- Signs and symptoms, a two-part notation



Writing the Diagnostic Statement

The diagnostic statement connects problem, etiology, and signs and symptoms. The first two parts of the statement are linked by "related to," some times abbreviated **R/T**. The last two parts are linked by "as evidenced by," some times abbreviated **AEB**.


E.G. *Ineffective Airway Clearance related to physiologic effects of pneumonia as evidenced by increased sputum, coughing, abnormal breath sounds, tachypnea, and dyspnea.*

- b) Collaborative problems – certain physiologic complications that nurse monitor to detect onset or changes in status. Nurses manage collaborative problems using physician – prescribed and nursing prescribed interventions to minimize the complications of the events.
3. Planning – development of goals and a plan of care designed to assist the patient in resolving the diagnosed problems. Setting

Establishing Expected Outcomes

An expected outcome is a measurable client behavior that indicates whether the person has achieved the expected benefit of nursing care. It may also be called a *goal* or *objective*. An expected outcome has the following characteristics:

- Client oriented
- Specific
- Reasonable
-




N.B. Dividing the nursing process into five distinct components or steps serves to emphasize the essential nursing actions that must be taken to resolve patient's nursing diagnoses and manage any collaborative problems or complications.

Critical thinking:

It is defined as an intellectually disciplined process of actively and skillfully conceptualizing, applying, analyzing, synthesizing, and or evaluating information gathered from, or generated by observation, experience, reflection, reasoning or communication, as a guide to belief and action. Critical thinking involves problem solving and decision making process, but it is a more complex process. Critical thinking competencies are the cognitive processes a nurse in clinical situations include diagnostic reasoning clinical inferences, and clinical decision-making. The nurse process is considered the specific critical thinking competency in nursing. Critical thinking skill assists the nurse to look at all aspects of a situation and then at a conclusion. When critical thinking is employed in clinical situations one should expect the how to examine ideas, beliefs, principles, assumptions, conclusions, statements and inferences before coming to a conclusion and make a decision

Study Questions:

1. Define nursing using modern definition.
2. List nurse professionals who significantly contributed to professional development.
3. State the historical trends of nursing development.
4. Mention steps in nursing process.
5. State two common ways of collecting data about client.
6. Describe critical thinking.



UNIT TWO

SAFETY IN HEALTH CARE FACILITIES

CHAPTER TWO

INFECTION CONTROL/ UNIVERSAL PRECAUTION

Learning Objectives:

At the end of this chapter the learner will be able to:

- Describe infection prevention in health care setups
- List chain of infection
- Identify between medical asepsis and surgical asepsis
- Discuss the purpose, use and components of standard precautions.
- Maintain both medical and surgical asepsis
- Describe how to setup a client's room for isolation, including appropriate barrier techniques.
- Identify hoe to follow specific airborne, droplet and contact precautions.

New Terminology

- | | |
|-----------------------|---------------------------------|
| - Airborne precaution | - protective isolation |
| - Contact precaution | - standard precaution |
| - Droplet precaution | - transmission-based precaution |
| - Isolation | |

Nursing Process

Assessment

- Identify appropriate times for hand washing
- Identify type of protective clothing required for barrier nursing.
- Identify epidemiology of the disease to determine how to prevent infection from spreading.
- Identify equipment needed to prevent spread of organisms
- Assess method of terminal cleaning and disposing equipment.
- Assess method of hand washing that is most appropriate for assigned task.
- Identify clients at risk for infection
- Assess availability of equipment for frequent hand washing.
- Evaluate health status of the nurse
- Check agency policy for hand washing protocol
- Assess need for use of unsterile gloves
- Assess nurses and clients for latex allergies
- Assess need for latex-free equipment and/or environment.

Planning/Objective

- To prevent the spread of endogenous and exogenous flora to other client.

- To reduce potential for transforming organisms from the hospital environment to the clients from acquiring nosocomial infections.
- To deliver client care with pathogen-free hands.
- To prevent pathogenic microorganisms spreading from client to client, environment or health care personnel to client.
- To prevent health care workers from contamination.

Implementation

- Preparation for isolation
- Donning and removing isolation attire
- Using a mask
- Assessing vital signs
- Removing items from isolation room
- Utilizing double-bagging for isolation
- Removing a specimen from isolation room
- Transporting isolation client outside the room
- Removing soiled large equipment from isolation room
- Hand washing (Medical asepsis)
- For using Waterless Antiseptic Agents
- Cleaning Washable Articles
- Donning (putting on) and Removing clean Gloves
- Managing Latex Allergies

Evaluation/Expected Outcome




Normal Body Defense

Individuals normally have defenses that protect the body from infection. These defenses can be categorized as non-specific and specific.

Specific Defenses

Specific defenses (immune): are directed against identifiable bacteria, viruses, fungi, or other infectious agents. Specific defenses of the body involve the immune system, which responds to foreign protein in the body (E.g. bacteria or transplanted tissues) or, in some



Inflammation is a local and non-specific defense response of the tissues to injury or infection. It is an adaptive mechanism that destroys or dilutes the injurious agent, prevents further spread injury, and promotes the repair of damage tissue. Inflammation is characterized by the following classic signs and symptoms of (Virchow, 1821-1902):

(a) Pain (dolor), (b) Swelling (tumor), (c) Redness (rubor), (d) Heat (calor), and (e) Impaired function of the part (functio laesa), if the injury is severe. Often words with “it is” describe an inflammatory process.

An infection is an invasion of the body tissue by microorganisms and their proliferation there. Such a micro organism is called infections agent.

Pathogenicity is the ability to produce disease; thus a pathogen is a microorganism that causes disease. A “true” pathogen causes disease or infection in a healthy individual. An opportunistic pathogen causes disease only in a susceptible individual. Etiology is the study of causes; the etiology of an infectious process is the identification of the invading microorganisms. Infectious diseases are the major cause of illness and death in Ethiopia.

Chain of

There are six steps of infection:

1. The micro organism
2. The place where the organism resides
3. Portal of exit
4. Method of transmission
5. Portal of entry
6. Susceptible host




Figure 1. Chain of infection

Conditions Predisposing to Infection

Certain conditions and invasive techniques predispose clients to infection because the integrity of the skin is broken or the illness itself establishes a climate favorable for the infectious process to occur. Among the most common are surgical wounds, changes in the antibacterial immune system, or alterations to the body.

Nosocomial Infection

Nosocomial infections are infections that are acquired while the client is in the hospital, infections that were not present or incubating at the time of admission.

Standard Precautions

Standard precaution is also called universal precautions. These were instituted as a result of the human immunodeficiency virus (HIV) epidemic. Blood and body fluid precautions were practiced on all clients regardless of their potential infectious state.

In 1987, body substance isolation (BSI) was proposed. The intent of this isolation system was to isolate all moist and potentially infectious body substances (blood, feces, urine, sputum, saliva, wound drainage and other body fluids) from all clients, regardless of their infectious status, primarily through the use of gloves.

Standard precaution blends the major features of universal precautions (blood and body fluids precautions) and body substance isolation into a single set of precautions to be used for the care of all

clients in hospitals, regardless of their diagnosis or presumed infection status. The new standard precautions apply to blood, all body fluids, secretions, and excretions, whether or not they contain visible blood; non-intact skin; and mucus membrane.

Fundamental Principles: Certain fundamental principles should be applied to all clients. These include hand washing, use of gloves, proper placement of clients in hospital to prevent spread of microorganisms to others or to the client, and appropriate use of isolation equipment to prevent the spread of microorganisms to health care workers and other clients.

Basic Medical Asepsis

Hand washing (Medical Asepsis)

Purpose

- To prevent the spread of infection
- To increase psychological comfort

Equipment

- Soap for routine hand washing
- Orange wood stick for cleaning nails, if available
- Running warm water, paper towel, trash basket

Procedure

1. Stand in front of but away from sink to avoid touching of uniform to a sink.
2. Ensure that paper towel is hanging down from dispenser.
3. Turn on water using foot pedal or faucet (using elbow of hand) so that flow is adequate, but not splashing.
4. Adjust temperature to warm. **Rationale:** cold does not facilitate sudsing and cleaning; hot is damaging to skin.
5. Wet hands under running water, wet hands facilitate distribution of soap over entire skin surface.
6. Place a small amount, one to two teaspoons (5-10mL) of liquid soap on hands. Thoroughly distribute overhands. Soap should come from a dispenser, possible; this prevents spread of microorganism.
7. Rub vigorously, using a firm, circular motion, while keeping your fingers pointed down, lower than wrists. Start with each finger, then between fingers, then palm and back of hand to create friction on all surfaces.
8. Wash your hands for at least 10-15 seconds. Duration of washing is important to produce mechanical action and allow antimicrobial products time to achieve desired effect.
9. Clean under your fingernails with an orangewood stick. (This should be done at least at start of day and if hands are heavily contaminated).

10. Rinse your hands under running water, keeping fingers pointed down ward in order to prevent contamination of arms.
11. Resoap your hands, rewash, and rerinse if heavily contaminated.
12. Dry hands thoroughly with a paper towel, while keeping hands positioned with fingers pointing up. Moist hands tend to gather more microorganisms from the environment.
13. Turn off water faucet with dry paper towel, if not using foot pedal to avoid contaminating the hands.
14. Restart procedure at step 5 if your hands touch the sink any time between steps 5 and 13.

Donning and Removing Gloves



3. Hold glove at wrist edge and slip finger into opening. Pull



- o Assess extent of barrier techniques needed (i.e., gloves, gown, mask, protective eye wear).
- o Assess need for special equipment (i.e., hazardous waste bags, plastic bags for specimens).

Planning

- o To prevent clients (especially compromised clients) from acquiring nosocomial infections.
- o To prevent the spread of micro organism to health professionals.
- o To reduce potential for transmission of micro organisms.
- o To protect hospital personnel and others from contamination.
- o To provide appropriate equipment and techniques for preventives measures.

Implementation/Procedures

Donning protective gear utilizing standard precautions.

Exiting a client's room utilizing standard precautions.

Evaluation/Expected outcomes

- Clients remain infection free.
- Transmission of micro organism is controlled.

- Health care workers protected from micro organism.
- Appropriate nursing interventions are carried out for the client.

Donning Protective Gear Utilizing Standard Precautions

Equipment

- Disposable gloves
- Gown
- Mask
- Apron
- Cap
- Protective eye wear (goggle)

Procedure

1. Wash hands using soap and dry.
2. Put on gown by placing one arm at a time though sleeves wrap gown around body so it cover clothing completely.
3. Bring waist ties from back to front of gown or turn back according to hospital policy. This ensures that entire clothing is covered by the gown, preventing accidental contamination.
4. Tie gown at neck or adhere velcro strap to gown

5. Don mask. Rationale: Masks are worn when there is an anticipated contact with respiratory droplet secretiveness.
6. Don protective eye wear such as face shield. Face shields will protect the nurse from splashing of blood or body fluids while caring for clients.
7. Done disposable gloves.

Standard Precaution Guidelines

- Wash hands thoroughly after removing gloves and before and after all client contact
- Wear gloves when there is direct contact with blood, body fluids, secretions, excretions, and contaminated items. This include neonate before first bath, wash as soon as possible if an anticipated contact with this body substances occurs.



- Place all contaminated articles and trash in leak proof bags.



Disinfection and sterilization

Disinfection: is a process that results in the destruction of most pathogens, but not necessarily their spores. Common methods of disinfection include the use of alcohol wipes, a hexachlorophene or chlorohexidine gluconate soap scrub, or povidone-iodine scrub to kill microorganisms on the skin.

Stronger disinfectants include phenol and mercury bichloride, which are too strong to be used on living tissue. Boiling can be used to disinfect inanimate objects. However, it does not destroy all microorganisms or spores.

Sterilization:

It is the process of exposing articles to steam heat under pressure or the chemical disinfectants long enough to kill all microorganisms and spores.

Exposure to steam at 18 pounds of pressure at a temperature of 125°C for 15 minutes will kill even the toughest organisms. A pressure steam sterilizer is called an autoclave.

Some chemicals also can be used to sterilize an object. However,

Items to be use to maintain sterility technique

Hair covering:

In sterile environments a cap or hood is worn to cover the hair. Remember that no hair can show. If the hair is long, a special type of hood will be worn.

Surgical Mask

In strict sterile situations such as in operation room (OR) or with protective isolation, the mask covers the mouth and nose. The purpose of mask is to form a barrier to stop the transmission of pathogens. In the OR or during other sterile procedure, the mask prevents harmful microorganisms in your respiratory tract from spreading to the client. When the client has an infection, the mask protects you from his/ her pathogens.

Sterile gown

Sterile gown is commonly worn in the OR, with protective isolation and some times in the delivery room. The hands touch only the part of sterile gown that will touch the


Sterile Gloves

For some procedures sterile gloves are worn. Remember that once gloves are put on, touching any thing unsterile contaminates them. Therefore, make all preparations before putting on gloves.

Procedures for putting on sterile gloves:

Steps:

1. Wash the hands to limit the spread of microorganisms
2. Open the outer glove package, on a clean, dry, flat surface



- b. Pull the second glove on; touching only the outside of the sterile glove with the other sterile gloved hand and keeping the fingers inside the cuff.
 - c. Adjust gloves and snap cuffs in to place. Avoid touching the inside glove and wrist area
- 8. Keep the sterile gloved hands above waist level. Make sure not to touch the cloths. Keep hands folded when not performing a procedure. Both actions help to prevent accidental contamination.



How ever, still some facilities follow such system.

Category-specific isolation, specific categories of isolation (eg. Respiratory, contact, enteric, strict or wound) are identified, using color-coded cards. This form of isolation is based on the client's diagnosis. The cards are posted outside the client's room and state that visitors must check with nurses before entering.

Disease –specific isolation, uses a single all-purpose sign. Nurse selects the items on the card that are appropriate for the specific disease that is causing isolation.

Preparing for Isolation

Purpose

To prevent spread of microorganisms

To control infectious diseases

Equipment

Specific equipment depends on isolation precaution system used.

- Soap and running water.
- Isolation cart containing masks, gowns, gloves, plastic bags isolation tape.
- Linen hamper and trash can, when needed.
- Paper towel
- Door card indicating precautions

Procedure

1. Check orders for isolation
2. Obtain isolation cart from central supply, if needed.
3. Check that all necessary equipment to carry out the isolation order is available.
4. Place isolation card on the client's door.
5. Ensure that linen hamper and trash cans are available, if needed.
6. Explain purpose of isolation to client and family.
7. Instruct family in procedures required.
8. Wash hands with antimicrobial soap* before and after entering isolation room.

* Types of antimicrobial soap or agent depend on infectious agent and client condition.

Donning and Removing Isolation Attire

Equipment

- Gown
- Clean gloves

Procedure

For donning attire

1. Wash and dry hands
2. Take gown from isolation cart or cupboard. Put on a new gown each time you enter an isolation room.
3. Hold gown so that opening is in back when you are wearing the gown.
4. Put gown on by placing one arm at a time through sleeves, put gown-up and over your shoulder
5. Wrap gown around your back, tying strings at your neck.
6. Wrap gown around your waist, making sure your back is completely covered. Tie string around your waist.
7. Don eye shield and/or mask, if indicated. Mask is required if there is a risk of splashing fluids.
8. Don clean gloves and pull gloves over gown wristlets.

For Removing Attire

1. Untie gown waist strings
2. Remove gloves and dispose of them in garbage bag.
3. Next, untie neck strings, bringing them around your shoulders, so that gown is partially off your shoulders.
4. Using your dominant hand and grasping clean part of wristlet, put sleeve wristlet over your non-dominant hand. Use your

non-dominant hand to up pull sleeve wristlet over your dominant hand.



5. Tie bottom strings around your neck to secure mask over your mouth. There should be no gaps between the mask and your face.
6. Important; change mask every 30 minutes or sooner if it becomes damp as effectiveness is greatly reduced after 30 minutes or if mask is moist.
7. Wash your hands before removing mask.
8. To remove mask, untie lower strings first, or slip elastic band off without touching mask.
9. Discard mask in a trash container
10. Wash your hands





Using Double-Bagging for Isolation

Equipment

2 isolation bags

Items to be removed from room

Gloves

Procedure

1. Follow dress protocol for entering isolation room, or, if you are already in the isolation room, continue with step 2.
2. Close isolation bag when it is one-half to three-fourths full. Close bag inside the isolation room.
3. Double-bag for safety if outside of bag is contaminated, if the bag could be easily penetrated, or if contaminated material in the bag is heavy and could break bag.
4. Set-up a new bag for continued use inside room. Bag is usually red with the word "Biohazard" written on outside of bag.
5. Place bag from inside room in to a bag held open by a second health care worker outside room if double bagging is required. Second health care worker makes a cuff with the top of the bag and places hands under cuff. This prevents hands from becoming contaminated.
6. Place bag in to second bag with out contaminating outside of bag. Secure top of bag by tying a knot in top of bag.


7. Take bag to designated area where biohazard material is



4. Help client to transport vehicle. Cover client with a bath blanket.
5. Tell receiving department what type of isolation client needs and what type of precaution hospital personnel should follow.
6. Remove bath blanket, and handle as contaminated linen when client returns to room.
7. Instruct all hospital personnel to wash their hands before they leave the area.
8. Wipe down transportation vehicle with antimicrobial solution if soiled.

Protocol for Leaving Isolation Room

Untie gown at wrist



Guide lines for Disposing of Contaminated Equipment

- Disposable gloves: place in isolation bag separate from burnable trash and direct to appropriate hospital area for disposal
- Glass equipment: Bag separately from metal equipment and return to CSR (Central Sterilization Room).
- Metal equipment: Bag all equipment together, label and return to CSR
- Rubber and plastic items: Bag items separately and return to CSR for gas sterilization.
- Dishes: Requires no special precautions unless contaminated with infected material; then bag, label and return to Kitchen.
- Plastic or paper dishes: Dispose of these items in burnable trash.
- Soiled linens: place in laundry bag, and send to separate area of laundry room from special care. If possible place linens hot-water-soluble bag. This method is safer for handling as bag may be placed directly into washing machine. (Double-bagging is usually required because these bags are easily punctured or torn. They also dissolve when wet.)
- Food and liquids: Dispose of these items by putting them in toilet – flush thoroughly.

-



CHAPTER THREE

CARE OF PATIENT UNIT

Learning Objective

At completion of this unit the learner will be able to:

- State the general instruction for nursing procedures.
- Define patient and patient unit.
- Take care of patient unit and equipment in health care facilities

General Instructions for all Nursing Procedures

1. Wash your hands before and after any procedure.




I. THE PATIENT UNIT

Definition:

Patient: A *Latin* word meaning to suffer or to bear.

- Is a person who is waiting for or undergoing medical/nursing treatment and care.

Patient Care Unit: is the space where the patient is accommodated



desired positions i.e. to elevate the head or the foot of the bed

- ⇒ Most commonly found in Ethiopia hospitals
- ⇒ Are less expensive and free of safety hazard
- ⇒ Handles should be positioned under the bed when not in use

C. Side rails

- It should be attached to both sides of the bed
 - _ Full rails – run the length of the bed
 - _ Half rails _ run only half the length of the bed and commonly attached to the pediatrics bed.

D. Bed Side Table/Cabinet

- Is a small cabinet that generally consists of a drawer and a cupboard area with shelves
- Used to store the utensils needed for clients care. Includes the washbasin (bath basin, emesis (kidney) basin, bed pan and urinal
- Has a towel rack on either sides or along the back
- Is best for storing personal items that are desired near by or that will be used frequently

E.g. soap, shampoo, lotion etc

E. Over Bed Table

- The height is adjustable

- Can be positioned and consists of a rectangular, flat surface supported by a side bar attached to a wide base on wheels
- Along side or over the bed or over a chair
- Used for holding the tray during meals, or care items when completing personal hygiene

F. *The Chair*

- Most basic care units have at least one chair located near the bedside
- For the use of the client, a visitor, or a care provider

G. *Overhead Light (examination light)*

- Is usually placed at the head of the bed, attached to either the wall or the ceiling
- A movable lamp may also be used
- Useful for the client for reading or doing close work
- Important for the nurse during assessment

H. *Suction and Oxygen Outlets*

- Suction is a vacuum created in a tube that is used to pull (evacuate) fluids from the body E.g. to clear respiratory mucus or fluids
- Oxygen is one of the gases frequently used for health care today. Oxygen is derived through a tube.

I. *Electrical Outlets*

- Almost always available in the wall at the head of the bed

J. *Sphygmomanometer*

- The blood pressure assessment tool, has two types:
 1. An aneroid
 2. Mercury, which is frequently used during nursing assessment.

K. *Call Light*

- Used for client's to maintain constant contact with care providers

II. *Care of Patient Unit*





- Protect table tops when using hot utensils or any solution that may leave stain or destroy the table top.
- Report promptly any damaged or missing equipment.

2. *Care of Equipment in General*

- Rinse used equipment in cold water. Soak materials in recommended antiseptic solutions. Remove any sticky material. Hot water coagulates the protein of organic material and tends to make it adhere.
- Wash well in hot soapy water. Use an abrasive, such as a stiff-bristled brush, to clean equipment.
- Rinse well under running water.
- Dry the article.
- Clean the gloves, brush and clean the sink.

3. *Care of Linen and Removal of Stains*

- Clean linen should be folded properly and be kept neatly in the linen cupboard.
- Dirty linen should be put in the dirty linen bag (hamper) and never be placed on the floor.
- Torn linen should be mended or sent to the sewing room.
- Linen with blood should be soaked in cold water to which a small amount of hydrogen peroxide is added if available.



UNIT THREE

BASIC CLIENT CARE

CHAPTER FOUR

ADMISSION, TRANSFER AND DISCHARGE OF PATIENTS

Learning Objectives:

At the end of this chapter students will be able to:

- Demonstrate how to orient a new client to the health care facility.
- Discuss concepts related to caring for the client's clothing and valuable items on admission.
- State some of the nursing consideration related to admission of a client.
- Demonstrate the ability to transfer a client from one unit to another safely and effectively.
- Identify nursing considerations related to a client's discharge from the health care facility.
- Explain teaching that should occur at the time of a client discharge.

A. Admission

Admission is a process of receiving a new patient to an individual unit (ward) of the hospital. (Hospitalized individuals have many needs and concerns that must be identified then prioritized and for which action must be taken).

Purpose

- To help a new patient to adjust to hospital environment and routines.
- To alleviate the patient's fear and worry about the hospitalization.
- To facilitate recovery of patient from his/her problems

Nurse's Responsibilities during Admission of a Patient to Hospital

1. Check for orders of admission.
2. Check about financial issue, payment scheme (free or paying)
3. Assess the patient's immediate need and take action to meet them. These needs can be physical (e.g. acute pain) or emotional distress, (upset)
4. Make introduction and orient the patient
 - Greet the patient
 - Introduce self to the patient and the family
 - Explain what will occur during the admission process (admission routines) such as admission bath, put on hospital gowns etc.


- Orient patient to individual unit: Bed, bathroom, call light, supplies and belonging; and how these items work for patient use.
 - Orient patient to the entire unit: location of nurses office, lounge etc.
 - Explain anything you expect a patient to do in detail. (This helps the patients participate in their care).
 - Introduce other staff and roommates.
4. Perform baseline assessment
- General assessment**
- a. Observation and physical examination such as:
- Vital signs; temperature pulse, respiration and blood pressure.
 - Intake and output
 - Measure the weight of the patient
 -

6. Documentation

- Record all parts of the admission process
- Other recording include
 - ⇐ Notification to dietary departments
 - ⇐ Starting kardex card and medication records
 - ⇐ If there is specific form to the facility, complete it.

N.B. Additional measures can be carried out according to the patient problems (diagnoses).

B. Transfer of the patient to another unit



- Ù Record the transfer in a transfer note. Give the time, the unit to which the transfer occurs, types of transportation (wheelchair, stretcher), and the client's physical and psychological condition
- Ù Make sure that the receiving unit is ready. Usually a short verbal report is given to the receiving department nurse.

C. Discharging a Patient

Indications for discharge

- Progress in the patient's condition
- No change in the patient's condition (Referral)
- Against medical advice
- Death

Nurse's Responsibility during Discharging a Patient

1. Check for orders that a patient need to be discharged
2. Plan for continuing care of the patient
 - Referral as necessary
 - Give information for a person involved in the patient care.
 - Contact family or significant others, if needed.
 - Facilitate transportation with responsible unit
3. Teaching the patient about
 - What to expect about disease outcome
 - Medications (Treatments)
 - Activity
 - Diet

- Need for continued health supervision, and others as needed
4. Do final assessment of physical and emotional status of the patient and the ability to continue own care.
 5. Check and return all patients' personal property (both items in patient unit and those kept in safe area).
 6. Help the patient or family to deal with business office for customary financial matters and in obtaining supplies.
 7. Keep records
 - o Write discharge note
 - o Keep special forms for facility

Discharge summaries usually include:

- Description of client's condition at discharge
- Treatment (e.g. Wound care, Current medication)
- Diet
- Activity level
- Restrictions

Referral is a condition in which a client/patient is sent to a higher health care system for better diagnostic and therapeutic actions.

- Any active health problems
- Current medication
- Current treatments that are to be continued
- Eating and sleeping habits

- Self-care abilities
- Support networks
- Life-style patterns
- Religious preferences

Discharging a patient against medical advice (AMA)

1. When the patient wants to leave an agency without the permission of the physician/nurse in charge – an authorized.
2. Ascertain why the person wants to leave the agency
3. Notify the physician/ nurse in charge of the client's decision
4. Offer the patient the appropriate form to complete
5. If the client refuses to sign the form, document the fact on the



Study Questions:

1. Mention concepts related to caring for clients belongings on admission.
2. State some of the nursing consideration related to admission of a client.
3. Exercise how to transfer a client from one unit to another safely and effectively.
4. Identify nursing considerations related to discharge of a client from health care facility
5. Explain teaching that should occur at time of a client discharge.



CHAPTER FIVE

VITAL SIGNS

Learning Objectives

At the end of the unit the learner will be able to:

- Describe the procedures used to assess the vital signs: temperature, pulse, respiration, and blood pressure.
- Identify factors that can influence each vital sign.
- Identify equipment routinely used to assess vital signs.
- Identify rationales for using different routes for assessing temperature.
- Identify the location of commonly assessed pulse sites.
- Take vital signs and interpret the finding.
- Document the vital signs.

II. Vital Signs (Cardinal Signs)

Vital signs reflect the body's physiologic status and provide information critical to evaluating homeostatic balance. The term "vital" is used because the information gathered is the clearest indicator of overall health status.

Vital sign Includes: T (temperature), PR (Pulse Rate), RR (Respiratory Rate), and BP (Blood Pressure)

Key Terminology

- apical pulse
- apex
- apnea
- axilla
- bradycardia
- bradypnea
- carotid pulse
- cheyne-stokes respiration
- cyanosis
- diastole
- dyspnea
- eupnea
- femoral pulse
- fever
- hypertension
- hypotension
- Korotokoff
- tympanic
- oral
- orthopnea
- palpation
- pedal pulse
- popliteal pulse
- pulse
- pulse pressure
- radial pulse
- rectal
- sphygmomanometer
- stethoscope
- systolic
- tachycardia
- temperature
- thermometer

Acronyms

BP	PR	PO	°C
RR	CVS	B	°F
T	O	BPM	

Purposes:

- To obtain base line data about the patient condition



4. Before and after the administration of certain medications that could affect RR or BP
(Respiratory and CVS (Cardio Vascular System))
5. Before and after surgery or an invasive diagnostic procedures
6. Before and after any nursing intervention that could affect the vital signs. E.g. Ambulation
7. According to hospital /other health institution policy.

I Temperature –Body temperature is the measurement of heat inside a person's body (core temperature); it is the balance between heat produced and heat lost.

Normal body temperature using oral (O; or per os, PO) measurement remains as appropriately 37⁰ celsius or 98.6 ⁰ F.

There are Two Kinds of Body Temperature

1. Core Temperature

- Is the Temperature of the deep tissues of the body, such as the cranium, thorax, abdominal cavity, and pelvic cavity
- Remains relatively constant
- Is the Temperature that we measure with thermometer

2. Surface Temperature:

- The temperature of the skin, the subcutaneous tissue and fat

Alterations in Body Temperature

Normal body temperature is 37⁰ C or 98.6 ⁰F (Average) the range is 36-38 ⁰c (96.8 – 100 ⁰F)

Pyrexia:



- Children's temperature continue to be more labile than those of adults until puberty
 - Elderly people, particularly those > 75 are at risk of hypothermia
 - Normal body temperature of the newborn if taken orally is 37 °C.
2. Diurnal variations (circadian rhythms)
- Body temperature varies through out the day
 - The point of highest body temperature is usually reached between 8:00 p.m. and midnight and lowest point is reached during sleep between 4:00 and 6:00 a.m.
3. Exercise
- Hard or strenuous exercise can increase body temperature to as high as 38.3 – 40 c – measured rectally
4. Hormones
- In women progesterone secretion at the time of ovulation raises body temperature by about 0.3 – 0.6°C above basal temperature.
5. Stress

- Stimulation of skin can increase the production of epinephrine and nor epinephrine – which increases metabolic activity and heat production.
6. Environment
- Extremes in temperature can affect a person's temperature regulatory systems.

Measuring Body Temperature

Sites to Measure Temperature

Most common are:

- Oral
- Rectal
- Axillary
- Tympanic

Thermometer: is an instrument used to measure body temperature

Types

1. Oral thermometer
 - Has long slender tips
2. Rectal thermometer
 - Short, rounded tips
3. Axillary
 - Long and slender tip
4. Tympanic

In other way it is also divided as mercury, digital and electronic types. In developed countries, mercury type thermometers are no more use in hospital setup but in our context still very important.

1. Rectal Temperature:

Readings are considered to be more accurate, most reliable, is $> 0.65^{\circ}\text{C}$ (1°F) higher than the oral temperature.

Procedure

- Explain the procedure to the patient
- Wash hands and assemble necessary equipment and bring to the patient bedside.
- Position the person laterally;
- Apply lubricant 2.5 cm above the bulb;
- Insert the thermometer 1.5 – 4 cm into the anus. For an infant 2.5cm, for a child 3.7 cm – for an adults 4 cm
- Measured for 2-3 minutes
- Remove the thermometer and read the finding
- Clean the thermometer with tissue paper
- A rectal thermometer record does not respond to changes in arterial temperature as quickly as an oral thermometer

Contraindications

- Rectal or perineal surgery;
- Fecal impaction – the depth of the thermometer insertion may be insufficient;

- Rectal infection;
- Neonates –can cause rectal perforation and ulceration;

2. Oral

Procedure

- Explain the procedure to the patient
- Wash hands and assemble necessary equipment and bring to the patient bedside.
- Position the person comfortably and request the patient to open the mouth;
- Hold the thermometer firmly with the thumb and fore finger; shake it with strong wrist movements until the mercury line falls to at least 35 °c.
- Place the bulb of the thermometer well under the client's tongue. Instruct the client to close the lips (not the teeth) around the bulb. Ensure that the bulb rests well under the tongue, where it will be in contact with blood vessels close to the surface.
- Remove the thermometer after 3 to 5 minutes, according to the agency guidelines.
- Remove the thermometer, wipe it using it once a firm twisting motion
- Hold the thermometer at eye level. Read to the nearest tenth



- Hold the glass thermometer in place for 8 to 10 minutes. Hold the electronic thermometer in place until the reading registers directly
- Remove and read the thermometer. Dispose of the equipment properly. Wash hands
- Record the reading

N.B. The axillary method is safest and most noninvasive.

Tympanic Temperature

The tympanic temperature is placed snugly in to the client's outer ear canal. It records temperature in 1 to 2 seconds. Many pediatric and intensive care units use this type of thermometer because it records a temperature so rapidly.

Procedure

- Wash the hands
- Explain the procedure to the client to ensure cooperation and understanding
-

child of 6 years or younger, use your nondominant hand to pull the ear down and back.

- Slowly advance the probe in to the client's ear with a back





Pulse Sites

Temporal: is superior (above) and lateral to (away from the midline of) the eye

1. **Carotid:** at the side of the neck below tube of the ear (where the carotid artery runs between the trachea and the sternocleidomastoid muscle)
2. **Temporal:** the pulse is taken at temporal bone area.
3. **Apical:** at the apex of the heart: routinely used for infant and children < 3 yrs

In adults – Left midclavicular line under the 4th, 5th, 6th intercostals space

Children < 4 yrs of the Lt. mid clavicular line

4. **Brachial:**



Method

Pulse: is commonly assessed by palpation (feeling) or auscultation (hearing)

The middle 3 fingertips are used with moderate pressure for palpation of all pulses except apical; the most distal parts are more sensitive,

Assess the Pulse for



Pulse Rate

-

Pulse Rhythm

- The pattern and interval between the beats, random, irregular beats – dysrhythmia

Pulse Volume: the force of blood with each beat

- A normal pulse can be felt with moderate pressure of the fingers and can be obliterated with greater pressure.
 - Full or bounding pulse forceful or full blood volume obliterated with difficulty
 - Weak, feeble or thready readily obliterated with pressure from the finger tips
 - Elasticity of arterial wall
 - A healthy, normal artery feels, straight, smooth, soft and pliable, easily bent after breaking
 - Reflects the status of the clients vascular system
- If the pulse is regular, measure (count) for 30 seconds and multiply by 2
- If it is irregular count for 1 full minute

Procedure for measuring radial pulse (the most common)

- ↳ Wash hands
- ↳ Explain the procedure to the client
- ↳ Position the client's fore arm comfortably with the wrist extended and the palm down
- ↳ Place the tips of your first, second, and third fingers over the client's radial artery on the inside of the wrist on the thumb side.

- ↳ Press gently against the client's radial artery to the point where pulsation can be felt distinctly
- ↳ Using a watch, count the pulse beats for 30 seconds and multiply by two to get the rate per minute
- ↳ Count the pulse for full minute if it is abnormal in any way or take an apical pulse
- ↳ Record the rate (BPM) on paper or the flow sheet. Report any irregular findings to appropriate person
- ↳ Wash your hands

III Respiration

Respiration is the act of breathing (includes intake of O_2 removal of CO_2)

Ventilation is another word, which refer to the movement of air in and out of the lungs.

Hyperventilation: very deep, rapid respiration

Hypoventilation: very shallow respiration

Two Types of Breathing

1. Costal (thoracic)
 - Involves the external muscles and other accessory muscles (sternocleidomastoid)
 - Observed by the movement of the chest up ward and down ward. Commonly used for adults

2. Diaphragmatic (abdominal)

-



Stress	"
Arteriosclerosis	"
Obesity	"
Hemorrhage	Decrease
Low hematocrit	"
External heat	"
Exposure to cold	Increase

Sites for Measuring Blood Pressure

1. Upper arm using brachial artery (commonest)
2. Thigh around popliteal artery
3. Fore -arm using radial artery



When taking blood pressure using stethoscope, the nurse identifies five phases in series of sounds called Korotkoff's sound.

Phase 1: The pressure level at which the 1st joint clear tapping sound is heard, these sounds gradually become more intense. To ensure that they are not extraneous sounds, the nurse should identify at least two consecutive tapping sounds.

Phase 2: The period during deflation when the sound has a swishing quality

Phase 3: The period during which the sounds are crisper and more intense

Phase 4: The time when the sounds become muffled and have a soft blowing quality

Phase 5: The pressure level when the sounds disappear

Procedure

Assessing Blood pressure

Purpose

- o To obtain base line measure of arterial blood pressure for subsequent evaluation
- o To determine the clients homodynamic status
- o To identify and monitor changes in blood pressure resulting from a disease process and medical therapy.

EQUIPMENT

- o Stethoscope
- o Blood pressure cuff of the appropriate size
- o Sphygmomanometer

Procedure


1. Prepare and position the patient appropriately
 - Make sure that the client has not smoked or ingested caffeine, within 30 minutes prior to measurement.
 - Position the patient in sitting position, unless otherwise specified. The arm should be slightly flexed with the palm of the hand facing up and the fore arm supported at heart level
 - Expose the upper arm
2. Wrap the deflated cuff evenly around the upper arm.
 - Apply the center of the bladder directly over the medial aspect of the arm. The bladder inside the cuff must be directly over the artery to be compressed if the reading is to be accurate.
 - For adult, place the lower border of the cuff approximately 2 cm above antecubital space.
3. For initial examination, perform preliminary palpatory determination of systolic pressure

- Palpate the brachial artery with the finger tips
 - Close the valve on the pump by turning the knob clockwise.
 - Pump up the cuff until you no longer feel the brachial pulse
 - Note the pressure on sphygmomanometer at which the pulse is no longer felt
 - Release the pressure completely in the cuff, and wait 1 to 2 minutes before making further measurement
4. Position the stethoscope appropriately
- Insert the ear attachments of the stethoscope in your ears so that they tilt slightly fore ward.
 - Place the diaphragm of the stethoscope over the brachial pulse; hold the diaphragm with the thumb and index finger.
5. Auscultate the client's blood pressure
- Pump up the cuff until the sphygmomanometer registers about 30 mm Hg above the point where the brachial pulse disappeared.
 - Release the valve on the cuff carefully so that the pressure decreases at the rate 2-3 mmHg per second.
 - As the pressure falls, identify the manometer reading at each of the five phases
 - Deflate the cuff rapidly and completely

- Repeat the above step once or twice as necessary to confirm the accuracy of the reading.
6. Remove the cuff from the client's arm
 7. For initial determination, repeat the procedure on the client's other arm, there should be a difference of no more than 5 to 10 mmHg between the arms. The arm found to have the higher pressure, should be used for subsequent examinations
 8. Document and report pertinent assessment data, report any significant change in client's blood pressure to the nurse in charge. Also report these finding:
 - A. Systolic blood pressure (of adult) above 140 mmHg.
 - B. Diastolic blood pressure (of an adult) above 90 mmHg
 - C. Systolic blood pressure of (an adult) below 100mmHg

Study questions

1. Explain vital signs and list what it includes.
2. Identify important times to assess vital signs.
3. Mention some of the factors affecting body temperature.
4. What does pulse deficit mean?
5. Define arterial blood pressure.
6. Explain the two methods of assessing blood pressure.



CHAPTER SIX

SPECIMEN COLLECTION

Learning Objectives:

At the end of this chapter, students will be able to:

- Identify at least three reasons for laboratory examination of urine.
- Demonstrate correct collection of the following urine specimens: midstream, 24-hours, fractional, and indwelling urine catheter.
- Explain at least one reason for collecting specimen like sputum, blood or stool.
- Demonstrate correct collection of a stool specimen.
- Demonstrate correct collection of a sputum specimen.

Key Terminology:

Hemoglobine

Hematocrite

Leukocyte

Occult



3. Assemble and organize all the necessary materials for the specimen collection.
4. Get the appropriate specimen container and it should be clearly labeled have tight cover to seal the content and placed in the plastic bag or racks, so that it protects the laboratory technician from contamination while handling it.
 - The patient's identification such as, name, age, card number, the ward and bed number (if in-patient).
 - The types of specimen and method used (if needed).
 - The time and date of the specimen collected.
6. Put the collected specimen into its container without contaminating outer parts of the container and its cover.

All the specimens should be sent promptly to the laboratory, so that the temperature and time changes do not alter the content.

A. Collecting Stool Specimen

Purpose

- For laboratory diagnosis, such as microscopic examination,

- Assist the patient and place the bed pan under the patient's buttocks (follow the steps under "Giving and removing bedpan")
- Give patient privacy by leaving alone, but not far
- Instruct the patient about how to notify you when finished defecation.
- Remove the bedpan and keep on safe place by covering it
- Recomfort the patient

3. Obtain stool sample

- Take the used bedpan to utility room/toilet container using spatula or applicator without contaminating the outside of the container.
- The amount of stool specimen to be taken depends on the purpose, but usually takes.
 - o 3.5 gm sample from formed stool
 - o 15.30 ml sample from liquid stool
- Visible mucus, pus or blood should be included into sample stool specimen taken.

4. Care of equipments and the specimen collected.

- Handle and label the specimen correctly
- Send the specimen to the laboratory immediately, unless there is an order for its handling. Because fresh specimen provides the most accurate results.

- Dispose the bedpan's content and give proper care of all equipments used.

5. Documentation and report

B. Collecting Urine Specimen

Types of urine specimen collection

1. Clean voided urine specimen
(Also called clean catch or midstream urine specimen)
2. Sterile urine specimen
3. Timed urine specimen
 - It is two types
Short period → 1-2 hours
Long period → 24 hours

Purpose

- For diagnostic purposes
 - Routine lab()s140tt

- Water and soap or cotton balls and antiseptic solutions (swabs).

For patients confined

- Urine receptacles (i.e. bedpan or urinals)
- Bed protecting materials
- Screen (if required)

Procedure

i) For ambulatory patients

Give adequate instruction to the patient about

- The purpose and method of taking specimen
- Assist the patient to move to the toilet

ii) For patient confined in bed

1. Prepare the patient unit providing privacy
2. Prepare the patient
 - Put on gloves
 - Place bed protecting materials under patient's hips
 - Assist the patient to position in bed and in p 517.re tF(ee tF(repa)81 TfF(e(Assi)9443.

3. Obtain urine specimen
 - Ask patient to void
 - Let the initial part of the voiding passed into the receptacle (bed pan or urinal) then pass the next part (the midstream) into the specimen container.
 - Hold the vulva or penis apart from the specimen container while the patient voids to decrease urine contamination.
 - Don't allow the container to touch body parts
 - Collect about 30-60 ml midstream urine
 - Handle the outside parts of the container and put on the cover tightly on specimen container



Collecting a Sterile Urine Specimen

Sterile urine specimen collected using a catheter in aseptic techniques (The whole discussion for this procedure presented on the catheterization part)

Collecting a Timed Urine Specimen

Purpose

- For some tests of renal functions and urine compositions, such as:- measuring the level of or hormones, such as adrenocortico steroid hormone creatinine clearance or protein quantitation tests.

Equipments Required

- Urine specimen collecting materials (usually obtained from the laboratory and kept in the patient's bathroom.)
- Format for recording the time, date started and end, and the amount of urine collected on each patient's voiding during the specified period for collection.

Procedure

1. Patient preparation

- Adequate explanation to the patient about the purpose of the test, when it begins and what to do with the urine
- Place alert signs about the specimen collection at the patient's bedside or bathroom.

- Label the specimen container to include date and time of each voiding as well as patient's identification data
- Containers may be numbered sequentially (e.g. 1st, 2nd, 3rd etc) in case of 24-hours urine collection.

2. Collecting the urine

- Usually it begin in the morning
- Before you begin the timing, the patient should void and do not use this urine (It is the urine that has been in the bladder some time)
- Then all urine voided during the specified time (e.g. the next 24 hours) is collected in the container
- At the end of the time (e.g. 24 hours period) the patient should void the last specimen, which is added to the rest.
- Ensure that urine is free of feces

C. Collecting sputum specimen

Sputum is the mucus secretion from the lungs, bronchi and trachea, but it is different from saliva. The best time for sputum specimen collection is in the mornings up on the patient's awaking (that have been accumulated during the night). If the patient fails to cough out, the nurse can obtain sputum specimen by aspirating pharyngeal secretion using suction.

Purpose

Sputum specimen usually collected for:

- Culture and sensitivity test (i.e. to identify the microorganisms and sensitive drugs for it)
- Cytological examination
- Acid fast bacillus (AFB) tests
-



- Put on gloves, to avoid contact with sputum particularly if hemoptysis (blood in sputum) present.
 - Ask pt to cough deeply to raise up sputum
 - Take usually about 15-30 ml sputum
 - Ask pt to spit out the sputum into the specimen container
 - Make sure it doesn't contaminate the outer part of the container. If contaminated clean (wash) with disinfectant
 - Cover the cap tightly on the container
3. Recomfort the patient
- Give oral care following sputum collection (To remove any unpleasant taste)
4. Care of the specimen and the equipments used
- Label the specimen container
 - Arrange or send the specimen promptly and immediately to laboratory.
 - Give proper care of equipments used
5. Document the amount, color, consistency of sputum, (thick, watery, tenacious) and presence of blood in the sputum.

D. Collecting Blood Specimen

The hospital laboratory technicians obtain most routine blood specimens. Venous blood is drawn for most tests, but arterial blood is drawn for blood gas measurements. However, in some setting nurses draw venous blood.

Purpose

Specimen of venous blood are taken for complete blood count, which includes

- Hemoglobin and hemotocrit measurements
- Erythrocytes (RBC) count
- Leukocytes (WBC) count
- Differential counts

Equipment

- Sterile gloves
- Tourniquet
- Antiseptic swabs
- Dry cotton (gauze)
- Needle and syringe
- Specimen container with the required diluting or preservative agents, for example: anticoagulant.
- Identification/ labeling: name, age address, etc.
- Laboratory requisition forms

Procedure

1. Patient preparation

- Instruct the pt what to expect and for fasting (if required)
 - Position the pt comfortably
2. Select and prepare the vein sites to be punctured
- Put on gloves
 - Select the vein to be punctured. Usually the large superficial veins used such as, brachial and median cubital veins.
 - Place the veins in dependent positions
 - Apply tourniquet firmly 15-20 cm about the selected sites. It must be tight enough to obstruct vein blood flow, but not to occlude arterial blood flow.
 - If the vein is not sufficiently to dilate massage (stroke) the vein from the distal towards the site or encourage the pt to clench and unclench repeatedly.
 - Clean the punctured site using antiseptic swabs
3. Obtain specimen of the venous to blood
- Adjust the syringe and needles
 - Clean/disinfect the area with alcohol swab, dry with sterile cotton swab
 - Puncture the vein sites
 - Release the tourniquet when you are sure in the vein
 - Withdraw the required amount of venous blood specimen

- Withdraw the needle and hold the sites with dry cotton (to apply pressure)
 - Put the blood into the specimen container
 - Made sure not to contaminate outer part of the container and not to distract the blood cells while putting it into the container
4. Recomfort the patient
5. Care of the specimen and the equipment
- Label the container
 - Shake gently (if indicated to mix)
 - Send immediately to laboratory, accompanying the request
 - Give care of used equipments
6. Documentation and reporting

Observations and Recording of Signs and Symptoms of the Patient


1. Objective Symptoms (signs):
- Are symptoms, which could be seen by the health personnel?
- E.g. swelling, redness, rash, body discharges (defecation, diaphoresis, emesis,)

2. Subjective Symptoms:

p Are symptoms, which are felt by the patient

E.g. decrease of appetite, dizziness, deafness, burning sensation, nausea, etc

3. Chart



- Accuracy
Records must be correct in all ways, be honest
- Completeness
No omission, avoid unnecessary words or statement
- Exactness
Do not use a word you are not sure of
- Objective information
Record what you see avoid saying (condition better)
- Legibility
Print/write plainly and distinctively as possible
- Neatness
No wrinkles, proper speaking of items
Place all abbreviation, and at end of statement
- Composition / arrangement
Chart carefully consult if in doubt avoid using of chemical formulas
- Sentences need to be complete and clear, avoid repetition
- Don't overwrite
- Don't leave empty spaces in between
- Time of charting
Specific time and date
- Color of ink
Black or blue (red for transfusion, days of surgery)

It should be recorded on the graphic sheet

All orders should be written and signed. Verbal or telephone orders should be taken only in emergency verbal orders should be written in the order sheet and signed on the next visit.

Orders of Assembling Patients Chart

- a. History sheet
 - b. Personal and social data
 - c. Order sheet
 - d. Doctor's progress notes
 - e. Nurses notes
 - f. Vital sign sheet (graphics)
 - g. Intake and output recording sheet
 - h. Laboratory and other diagnostic reports
 - Patients or relatives and friends of patients are not allowed to read the chart when necessary but can have access if allowed by patient.
4. Intake and out put
- a. Intake: all fluids that is taken in to the body through the mouth, NG tube or parentrally
 - b. Output: all fluid that is excreted or put out of the body through the mouth. N/G tube, urethra, drainage tube or other route (GI-diarrhea, vomiting).

Purpose:

- To replace fluid losses

- To provide maintenance requirements
- To check for retention of body fluid

Fluid balance sheet

- ◆ 24 hrs the intake out put should be compared and the balance is recorded

 p Positive balance if intake > output

Negative balance if out put > intake

Study Questions

1. Explain at least three reasons for laboratory examination of urine.
2. Explain at least one reason for collecting specimens like sputum, blood or stool.
3. Mention purposes for sputum specimen collection.
4. Describe the process how to draw venous blood for laboratory investigation.
5. How can you obtain sterile urine specimen?
6. Differentiate between signs and symptoms.



CHAPTER SEVEN

BED MAKING

Learning Objectives

At the end of this unit, the learner able to:

- Describe different types of bed making.
- State the purposes of bed making in health care facilities.
- Develop understanding about general instruction of bed making
- Develop a skill to make different types of bed.
- Explain the purposes of side rails.
- List necessary equipment for bed making.
- Arrange bed-making equipment in order of their use.

Key terminology



Anesthetic bed: is a bed prepared for a patient recovering from anesthesia

⇒ **Purpose:** to facilitate easy transfer of the patient from stretcher to bed

Amputation bed: a regular bed with a bed cradle and sand bags

⇒ **Purpose:** to leave the amputated part easy for observation

Fracture bed: a bed board under normal bed and cradle

⇒ **Purpose:** to provide a flat, unyielding surface to support a fracture part

Cardiac bed: is one prepared for a patient with heart problem

⇒ **Purpose:** to ease difficulty in breathing

General Instructions

1. Put bed coverings in order of use
 2. Wash hands thoroughly after handling a patient's bed linen
- Linens and equipment soiled with secretions and excretions harbor micro-organisms that can be transmitted directly or by

8. Bed spread

Note

- *Pillow should not be used for babies*
- *The mattress should be turned as often as necessary to prevent sagging, which will cause discomfort to the patient.*

A. Closed Bed

- It is a smooth, comfortable, and clean bed that is prepared for a new patient

Essential Equipment:

- Two large sheets
- Rubber draw sheet
- Draw sheet
- Blankets
- Pillow cases
- Bed spread

Procedure:

- Wash hands and collect necessary materials
- Place the materials to be used on the chair. Turn mattress and arrange evenly on the bed
- Place bottom sheet with correct side up, center of sheet on center of bed and then at the head of the bed

- Tuck sheet under mattress at the head of bed and miter the corner
- Remain on one side of bed until you have completed making





- Never turn a helpless patient away from you, as this may cause him/her to fall out bed
- When you have made the patient comfortable and secure as near to the edge of the bed as possible, to go the other side carrying your equipment with you
- Loosen the bedding on that side
- Fold, the bed spread half way down from the head
- Fold the bedding neatly up over patient
- Roll dirty bottom sheet close to patient
- Put on clean bottom sheet on used top sheet center, fold at center of bed, rolling the top half close to the patient, tucking top and bottom ends tightly and mitering the corner
- Put on rubber sheet and draw sheet if needed
- Turn patient towards you on to the clean sheets and make comfortable on the edge of bed
- Go to the opposite side of bed. Taking basin and wash cloths with you, give patient back care
- Remove dirty sheet gently and place in dirty pillow case, but not on the floor
- Remove dirty bottom sheet and unroll clean linen
- Tuck in tightly at ends and miter corners
- Turn patient and make position comfortable
- Back rub should be given before the patient is turned on his /her back
- Place clean sheet over top sheet and ask the patient to hold it if she/he is conscious

- Go to foot of bed and pull the dirty top sheet out
- Replace the blanket and bed spread
- Miter the corners
- Tuck in along sides for low beds
- Leave sides hanging on high beds
- Turn the top of the bed spread under the blanket
- Turn top sheet back over the blanket and bed spread
- Change pillowcase, lift patient's head to replace pillow.
Loosen top bedding over patient's toes and chest
- Be sure the patient is comfortable
- Clean bedside table
- Remove dirty linen, leaving room in order
Wash hands

Bed Making

Making a post operative bed

- o The entire bed need clean linen.
- o Make the bottom of bed as you normally would. The post operative the bottom of bed as you normally would. The post operative bed usually requires a draw sheet under the client's hips. Usually another draw sheet is placed under the client's head.
- o In some cases, top liners are simply tan-folded to the foot of the bed. In others, a full post operative bed is made.

To do this, put the top linens over the foundation, but do not tuck them in. Fold down the top as you would do in an occupied bed. Then fold the bottom of the linens up so that the fold is even with the bottom of the mattress. Do not tuck the linen in. Fanfold the top linens to the side so that they lay opposite from where you will place the client's stretcher. Alternatively, you may fanfold the linens to the foot of the bed. Leave a tab on top for easy grasping.

- o Have two or more pillows available, but do not put them on the bed. Rational: A pillow may be contraindicated for a client, usually the physician or charge nurse will determine when it is safe for the client to have one.
- o Be sure all furniture is out of the way.
- o Be sure the call light is available, but keep it on the bed side stand until the client is in bed.



re

N.B. Procedures for other beds like cardiac bed are similar except the following points.

- ↳ For cardiac patient the bed need extra materials such as over bed table and additional pillows
- ↳ Hard board is needed under the mattress for fracture bed.

Study questions

1. How many types of bed making do you know?
2. What is the function of bed the cradle?
3. Which types of bed are usually prepared for newly admitted patients?
4. What is the difference between open and closed bed?
5. Define occupied bed.



CHAPTER EIGHT

PERSONAL HYGIENE AND SKIN CARE

Learning Objectives:

At the end of this chapter the learners will be able to:

- State the purposes of giving mouth care
- Demonstrate the skill of assisting a client with oral care
- Demonstrate for cleansing and caring for dentures
- Demonstrate caring for client's fingernails, and toes nails, addressing reasons for attention of each other.
- List reasons for routine hair care
- Describe and demonstrate giving a backrub, hand and foot massage, and foot soak
- Demonstrate how to assist a client with cleansing bath.

Key terminology

Halitosis

Nits

Pediculosis

Perineal care

A. Mouth Care

Purpose

- To remove food particles from around and between the teeth
- To remove dental plaque to prevent dental caries
- To increase appetite
- To enhance the client's feelings of well-being
- To prevent sores and infections of the oral tissue
- To prevent bad odor or halitosis

Equipments


- Toothbrush (use the person's private item. If patient has none use of cotton tipped applicator and plain water)
- Tooth paste (use the person's private item. If patient has none of use cotton tipped applicator and plain water)
- Cup of water
- Emesis basin
- Towel
- Denture bowl (if required)
- Cotton tipped applicator, padded applicator
- Vaseline if necessary

Procedure

1. Prepare the pt:
 - Explain the procedure

- Assist the patient to a sitting position in bed (if the health condition permits). If not assist the patient to side lying with the head on pillows.
 - Place the towel under the pt's chin.
 - If pt confined in bed, place the basin under the pt's chin
2. Brush the teeth

-



2. To floss the upper teeth. Use the thumb and index finger to stretch the floss. Move the floss up and down between the teeth from the tops of the crowns to the gum
3. To floss the lower teeth, use your index fingers to stretch the floss

Note: *If the patient has denture, remove them before starting and wash them with brush*

Mouth care for unconscious patient

Position

- Side lying with the head of the bed lowered, the saliva automatically runs out by gravity rather than being aspirated by the lungs or if patient's head can not be lowered, turn it to one side: the fluid will readily run out of the mouth, where it can be suctioned
- Rinse the patient's mouth by drawing about 10 ml of water or mouth wash in to the syringe and injecting it gently in to each side of the mouth
- If injected with force, some of it may flow down the clients throat and be aspirated into the lung
- All the rinse solution should return; if not suction the fluid to prevent aspiration

Giving and Receiving Bedpans and Urinals

- Bedpan is a material used to receive urine and feces in females and feces in male

- Urinal -is used to receive urine
- Are of two types male and female

Types of Bedpan

1. The high back, or regular pan (standard pan)
2. A fracture, the slipper or low back pan

Advantage

- ⇒ Has a thinner rim than as standard bed pan
- ⇒ Is designed to be easily placed under a person's buttocks

Disadvantage

- ⇒ Easier to spill the contents of the fracture pan
- ⇒ Are useful for people who are
 - a. Paralyzed or who cannot be turned safely (e.g. Spinal injury)
 - b. Confined in a body or long leg cast
 - c. Immobilized by some types of fracture
 - d. Very thin or emaciated

3. The pediatric bedpan
 - Are small sized
 - Usually made of a plastic

B. Bath (Bathing and Skin Care)

It is a bath or wash given to a patient in the bed who is unable to care for himself/herself.

1. Cleansing bath: Is given chiefly for cleansing or hygiene purposes and includes:

- Complete bed bath: the nurse washes the entire body of a dependent patient in bed
- Self-help bed bath: clients confined to bed are able to bath themselves with help from the nurse for washing the back and perhaps the face
- Partial bath (abbreviated bath): only the parts of the client's body that might cause discomfort or odor, if neglected are washed the face, hands, axilla, perineum and back (the nurse can assist by washing the back) omitted are the arms, chest, and abdomen.
- Tub bath: preferred to bed baths because it is easier to wash and rinse in a tub. Also used for therapeutic baths
- Shower: many ambulatory clients are able to use shower
- The water should feel comfortably warm for the client
- People vary in their sensitivity to heat generally it should be 43-46 °c (110-115°f)
- The water for a bed bath should be changed at least once


Before bathing a patient, determine

- a. The type of bath the client needs
- b. What assistance the client needs

- c. Other care the client is receiving – to prevent undue fatigue
- d. The bed linen required

Note: *when bathing a client with infection, the caregiver should wear gloves in the presence of body fluids or open lesion.*

Principles





- Basin with warm water (43-46⁰c for adult and 38-40⁰c for children)
- Soap on a soap dish
- Hygienic supplies, such as, lotion, powder or deodorants (if required)
- Screen
- Disposable gloves
- Lotion thermometer (if available)

Procedures

1. Prepare the patient unit
 - Close windows and doors, use screen to provide privacy.
2. Prepare the patient and the bed
 - Place the bed in high position to reduce undue strain on the nurse's back
 - Remove pt's gown and pajamas
 - Assist pt to move toward you so it facilitates access to reach pt without undue straining. Position the pt in supine, semi -Fowler's or Fowler's depending on the pt's condition.

Check the temperature of the water using lotion thermometer /back of the hand.

3. Make a bath with the washcloth, so it retains water and heat than a cloth loosely held

4. Washing body parts
 - Expose only the parts of the patient's body being washed avoid unnecessary exposing.
 -




- i. Give proper care of materials used for bathing
 - Document and report pertinent data
 - Observation of the skin condition
 - General appearance or reaction of the pt
 - Type of bath give

Report any abnormal findings to the nurse in charge

B. Therapeutic Baths

- Are usually ordered by a physician/ nurse



- [EPHTI Basic Clinical Nursing Skills](#) (2014)



- Clean bed linen
- Bath thermometer if available
- Disinfectant for cleansing the tub

Procedure

Ù



- ↳ Don't leave a child or a client who is unsure, unsteady, self injurious alone.
- ↳ When the client has finished bathing, help the client out of tub and to dry. After dressing assist the client back to the room
- ↳ Inform the cleaner to carefully clean the tub after the bath
- ↳ Dispose of the glove and wash your hands
- ↳ Document the procedure, describing any unusual client reactions

Back Care (massage): includes the area from the back and shoulder to the lower buttocks

Purpose

- To relieve muscle tension
- To promote physical and mental relaxation
- To improve muscle and skin functioning
- To relieve insomnia
- To relax patient
- To provide a relieve from pain
- To prevent pressure sores (decubitus)
- To enhance circulation

Equipment

Basin of warm water
Washcloth
Towel
Soap

Skin care lotion

Procedure

1. Prepare the pt and pt's unit
 - Provide privacy by using screen or closing windows and doors.
 - Assist pt to move close to your working side
 - Position patient prone (lie on abdomen) if possible. If not because of the pt's condition, use side lying position with the pt facing away from you.
 - Expose the back of the pt.
 - Spread towel close to pt's back to protect foundation of the bed.
 - Wash the back with warm water and soap using wash

- Pause at the neckline, using your fingers to massage the side of the neck.
-



- Leave the pt comfortably
4. Give proper care of equipments
 5. Document the procedure, your observations and pt's reactions
 - Report any abnormal observations on the skin of the back (such as signs of pressure sore) to the nurse and physician in charge of the pt.

Three Types of Massage Strokes

1. Effleurage: stroking the body
2. Light, circular friction and straight, deep, firm, strokes
3. Petrissage: kneading and making large quick pinches of the skin, tissue, and muscle
 - Clean the back first
 - Warm the massage lotion or oil before use by pouring over your hands: cold lotion may startle the client and increase discomfort
 1. Effleurage the entire back: has a relaxing sedative effect if slow movement and light pressure are used
 2. Petrissage first up the vertebral column and then over the entire back: is stimulating if done quickly with firm p

- Assess: signs of relaxation and /or decreased pain (relaxed breathing, decreased muscles tension, drowsiness, and peaceful affect)
 - ⇒ Verbalizations of freedom from pain and tension
 - ⇒ Areas of redness, broken skin, bruises, or other signs of skin breakdown

Note

- *The duration of a massage ranges from 5-20 minutes*
- *Remember the location of bony prominence to avoid direct pressure over this areas*
- *Frequent positioning is preferable to back massage as massaging the back could possibly lead to subcutaneous tissue degeneration.*

NB. Backrub requires special skills as it might cause subcutaneous tissue degeneration; mainly in elderly.





For the client who has flatulence: limit carbonated beverages; avoid gas-forming foods

4. Exercise

- Regular exercise helps clients develop a regular defecation pattern and normal feces

5. Positioning

- Sitting position is preferred

Measures to assist the person to void include:

- Running water in the sink so that the client can hear it
- Warming the bed pan before use
- Pouring warm water over the perineum slowly
- Having the person assume a comfortable position by raising the head of the bed (men often prefer to stand)
- Providing sufficient analgesia for pain
- Having the person blow through a straw into a glass of water – relaxes the urinary sphincter

Perineal Care (Perineal – Genital Care)

Perineal Area:

- Is located between the thighs and extends from the symphysis pubis of the pelvic bone (anterior) to the anus (posterior).
- Contains sensitive anatomic structures related to sexuality, elimination and reproduction



Equipments

- Bath towel
- Cotton balls and gauze squares
- Pitcher with warm water or/and prescribed solution in container
- Gloves
- Bed pan
- Bed protecting materials
- Perineal pad or dressing (if needed)

Procedure

1. Patient preparation
 - Give adequate explanation
 - Provide privacy
 - Fold the top bedding and pajamas (given to expose

For Female

- Remove dressing or pad used
- Inspect the perineal area for inflammation excoriation,



Care

- Convenient for a woman to be on a bed pan to clean and rinse the vulva and perineum
- Secretion collects on the inner surface of the labia
- Use one hand to gently retract the labia
- Use a separate section of wash cloth for each wipe in a downward motion (from urethra to back perineum)
- Then clean the rectal area

Note

- *Following genital or rectal surgery, sterile supplies may be required for cleaning the operative site, E.g. Sterile cotton balls*
- *The operative site and perineal area may be washed with an antiseptic solution – apply by squirting them on the perineum from a squeeze bottle*

Male Perineum

- The penis contains pathways for urination and ejaculation through the urethral orifice (meatus)
- At the end of the penis is the glans covered by a skin flap (fore skin or prepuce)
- The urethral orifice is located in the center of the penis and opens at the tip



Combing/Brushing of Hair

A patient hair should be combed and brushed daily most patients do this themselves if the required materials provided and others may need nurse's help (assistance)

Purpose

- Stimulates the blood circulation to the scalp
- Distribute hair oils evenly and provide a healthy sheen
- Increase the patient's sense of well-being.

Equipments

- Comb (which is large with open and long toothed)
- Hand mirror
- Towel
- Lubricant/oils (if required)

Procedure

1. Prepare the patient
 - Position the patient in either sitting or semi-fowler's or flat, if the pt is weak to seat or unconscious.
 - Place the towel over the patient's shoulder, if in sitting position or over the pillow if pt is in semi-fowler or lying position.
 - Remove any pins and ribbons
2. Comb the hair by dividing the hair

- Hold a section of hair 2-3 inches from the end and comb the end until it is free from tangles. Move towards the scalp by combing in the same manner to remove tangles.
 - Continue fluffing the hair outward and upward until all the hairs combed.
 - Arrange the hair as neatly and simply as possible according to the patient's preference of style.
3. Recomfort the pt
- Remove the towel
 - Put patient in comfortable position
4. Care of equipment
5. Documentation

E. Shampooing/Washing the Hair of Patient Confined to Bed

Purpose

- Stimulate blood circulation to the scalp through massaging
- Clean the patients hair so it increase a sense of well-being to the pt
- To treat hair disorders like dandruff

Equipments

- Comb and brush
- Shampoo/soap in a dish

- Shampoo basin
- Plastic sheet
- Two wash towels
- Cotton balls
- Water in basin and pitcher
- Receptacle (bucket) to receive the used water
- Lubricants/oil as required

Procedure

1. Prepare the patient
 - Assist patient to move to the working side of the bed
 - Remove any hair accessories (e.g. pins, ribbons etc)
 - Brush and comb the hair to remove tangles
2. Arrange the equipments
 - Place the plastic sheet under patient's head and shoulder
 - Remove the pillows from under the pt's head and place it under pt's shoulder (to hyper extend the neck)
 - Tuck the towel under the pt's shoulder and neck
 - Place (arrange) the shampoo basin under the pt's head with one end extending to the receptacle for used water.
 - If there is no shampoo basin, use the plastic sheet, which is under pt's shoulder and head, make a funnel type fold and extend it to the receptacle.

-



6. Ensure pt's comfort
 - Remove plastic sheet shampoo basin
 - Assist pt for comfortable position
 - Assist pt in grooming

7. Care of equipment

8. Documentation and reporting

Pediculosis Treatment

Definition

Pediculosis: infestation with lice

Purpose

- To prevent transmission of some arthropod born diseases
- To make patient comfortable

Equipment

Lindane

1% permethrine cream rinse

Clean linen

Fine-tooth "nit" comb

Disinfectant for comb


Clean gloves

Towel

Lice:

- Are small, grayish white, parasitic insects that infest mammals
- Are of three common kinds:

- b Pediculose capitis: is found on the scalp and tends to stay hidden in the hairs
- b Pediculose pubis: stay in pubic hair
- b Pediculose corporis: tends to cling to clothing, suck blood from the person and lay their eggs the clothing suspect their presence in the cloth and the body:
 - a. The person habitually scratches





- 1.5% solution of Gammaxine effective to kill the adult lice in one application
- Does not kill nits
- Should be repeated to kill the newly hatched nits, for complete elimination
- The lotion is applied over scalp after a clean soapy wash of hair
- After 12-24 hrs the scalp is washed with soap to remove the lotion
- Avoid contact with lice
- Can also be used for pubic and body lice

F. Feeding a Helpless Patient

During illness, trauma or wound healing, the body needs more nutrients than usual. However, many peoples, because of weakness, immobility and/or one or both upper extremities are unable to feed themselves all or parts of the meal. Therefore, the nurse must be knowledgeable, sensitive and skillful in carrying out feeding procedures.

Purpose

- To be sure the pt receives adequate nutrition
- To promote the pt well-beings

Procedure

1. Prepare pt units

- Remove all unsightly equipments; remove solid linens and arranging bedside tables.
- Control unpleasant odors in the room by refreshing the room. Odor free environment makes eating more pleasant and aids digestion.

2. Prepare the patients

- Offers bedpan and urinals. To comfort pt and avoid interruption by elimination needs.
- Assist pt to wash hands, face and oral care
- Position patient comfortably
 - ⇒ Mid or high Fowler's position
- Protect the bed using suitable protective cover

3. Prepare the food



Consists of:

- Providing a urinal or bed pan if client is confined to bed
- Washing the face and hands and
- Giving oral care

Late Morning Care

- Is provided after clients have breakfast

Includes:

- The provision of a urinal or bed pan
- A bath or shower
- Perineal care
- Back massage and
- Oral, nail and hair care
- Making clients bed

Afternoon Care

- When clients return from physiotherapy or diagnostic tests
- Includes:
 - ↳ Providing bed pan or urinal
 - ↳ Washing the hands and face
 - ↳ Assisting with oral care refresh clients

Evening Care

- Is provided to clients before they retire for the night




CHAPTER NINE

COLD AND HEAT APPLICATION

Learning Objectives:

At the end of this chapter the learners will be able to:



- Part of alcohol to 3 parts of Luke warm H₂O remove patient's gown
- Take the patient temperature, sponge the body using the wash cloth alternately, sponge each part 2-3 min. changing the was cloth
- Heat loss is by conduction or vaporization
- Check pulse frequently and report any change

Local Application of Heat and Cold

Heat and cold are applied to the body for local and systemic effects

Heat Application

Purpose

1. To relieve pain and muscles spasm – by relaxing muscles
 - Increase blood flow to the area
2. To relieve swelling (facilitate wound healing)
 - To relieve inflammation and congestion

Heat

- Increases the action of phagocytic cells that ingest moisture and other foreign material
 - Increases the removal of waste products or infection metabolic process
3. To relieve chilling and give comfort
- Heat can be applied in both dry and moist forms

Dry Heat :- is applied locally, for heat conduction

- By means of a hot water bottle



The temperature of the water is 32 °C (below body temperature) 27-37 – alcohol evaporates at a low temperature and therefore removes body heat rapidly

- Less frequently used – because alcohol causes skin drying
- Heat loss is by conduction and vaporization
- Determine the patients' temperature, PR and RR frequently every (Q) 15 min
- Sponge each area (part) for 2-3 min changing the wash cloth
- The sponge bath should take about 30 minutes
- Reassess v/s at the end
- Discontinue the bath if the clients becomes pale or cyanotic or shivers, or if the PR becomes rapid or irregular

Temperature of hot water bottle (bag) 52 °C for normal adults, 40.5 – 46 °C – for debilitated (unconscious patients).

40.5-46 °C for children < 2 yrs;

Fill the bag about 2/3 full;

Expel the remaining air and secure the top;

Maximum effect occurs in 20-30 min;

The application is repeated Q2 – 3 hrs to relieve swelling compress – a moist gauze or cloth immersed in (hot or cold) water and applied over an area.

Local Application of Cold and Heat

Application of Cold

- Has systemic and local effect
- Can be applied to the body in two ways
 1. Moist
 2. Dry

Purpose: (Indication)

- To reduce body temperature during high fever and hyper pyrexia or sun stroke
- To relieve local pain
- To reduce subcutaneous bleeding e.g. in sprain and contusion
- To control bleeding e.g. epistaxis
- To relieve headache
- To provide comfort to a patient in extreme hot weather if desired

1. Moist Cold

- Cold compress
- A cloth (padded gauze) is immersed in cold water and applied in area where we get large superficial vessels
E.g. axilla and groin
- Change the cloth when it becomes warm
- Applied for 15-20 min

2. Dry Cold (Ice Bag)

- Ice kept in a bag
- Covered with cloth and applied on an area
- Temperature $<15^{\circ}\text{C}$

Application of Heat

Purpose

- To relieve stasis of blood
- To increase absorption of inflammatory products
- To relieve stiffness of muscle and muscle pain
- To relieve pain and swelling of a localized inflammation boil or carbuncle – sometimes increases edema, increases capillary permeability
- To increase blood circulation
- To promote suppuration
- To relieve distention and congestion
- To provide warmth to the body

Methods

1. Dry Heat

- Using hot water bottle (bags)
- After contact of the body with moisture of water vapors temperature $>46^{\circ}\text{C}$
- 52°C for normal adults

- 40.5 – 46°C for debilitated or unconscious patient's and child < 2 yrs
- 2/3 of the bag should be filled with water
- Expel the remaining air and secure the top
- Dry the bag and hold it upside down to test for leakage
- Wrap it in a towel or cover and place it on the body part
- Maximum effect occurs in 20-30 min
- Remove after 30-45 minutes

2. **Moist Heat**

1. Hot compress: a wash cloth immersed in hot water of temperature 40-46°C and change the site of washcloth frequently

Complication

- Paralysis
- Numbness
- Loss of sensation – fear of burn

2. Sitz bath

Sitz Bath (hit bath)

It is used to sock the client's pelvic area

- A clients sits in a special tub or a bowel
- The area from the mid thighs to the iliac crests or umbilicus - increases circulation to the perineum (when the legs are also immersed blood circulation to the perineum or pelvic area decrease)

- Temperature of water – 40-43 °c (105-110 ° F) – unless the patient is unable to tolerate the temperature

Purpose:

- To relieve pain in post operative rectal condition
- Smoothen irritated skin (perineum)
- Facilitates wound healing (after episiotomy)
- To release the bladder in case of urinary retention

If it is going to be given in the tub – fill ½ the tube with water and add the ordered medication

In a bowel – fill 2/3 of it with water – add the ordered medication and dilute

The medication to Rx the perineum in KMNO₄ sol. 250 mg KMNO₄ in 500 ml of water

The duration of the bath is

Study questions:

1. Mention the two purposes of the heat application.
2. Describe the mechanism of action of heat application to effect its purposes.
3. What is tepid sponge?
4. What is the common medicine used in sitz bath?
5. What is the average duration of time the patient is soaked in sitz bath?



CHAPTER TEN

BODY MECHANICS AND MOBILITY

Learning Objectives:

At the end of this chapter the learner will be able to:

- State the principle underlying proper body mechanics and relate a nursing consideration.
- State the purposes of range of motion exercise.
- Identify principles related to safe movement of clients in and out of bed.
- Demonstrate the ability to move a partially mobile client safely from bed to chair and back.
- Demonstrate the ability to teach each of the crutch walking gaits to a client.
- Mention different positions used for various examination and treatment.

Key Terminology

Body alignment

Dorsal lithotomy

Prone

Base of support	Foot drop
Protective device	
Body mechanics	Fowler's position
Recumbent	
Center of gravity	Gait
Rotation	
Contracture	Gaitbelt
Transfer belt	
Centrolateral	Line of gravity
Sim's position	
Dangling	Paralysis
Supnation	

Acronyms

AROM
PROM
ROM

Body Mechanics: is the effort; coordinated, and safe use of the body to produce motion and maintain balance during activity.

Proper Body Mechanics


Use of safest and most efficient methods of moving and lifting is called body mechanics. This means applying mechanical principles of movements to the human body.

Basic Principles of Body Mechanics

The laws of physics govern all movements. From these laws we derive the general principles of body mechanics.

Basic Principles

1. It is easier to pull, push, or roll an object than to lift it. The movement should be smooth and continuous, rather than jerky.



area, half below it, when thinking of the body divided horizontally. In addition, half the body weight is to each side, when thinking the body divided vertically. When lifting an object, bend at knees and hips, and keep the back straight. By doing so, the center of gravity remains over the feet, giving extra stability. It is thus easier to maintain balance.

Base of Support

A person's feet provide the base of support. The wider the base of support, the more stable the object within limits. The feet are spread side wise when lifting, to give side-to-side stability. One foot is placed slightly in front of the other for back-to-front stability. The weight is distributed evenly between both feet. The knees are flexed slightly to absorb jolts. The feet are moved to turn the object being moved.

Line of Gravity

Draw an imaginary vertical (up and down) line through the top of the head, the center of gravity, and the base of support. This becomes



- o To increase the patient sense of independence and self-esteem
- o To assist a patient who is unable to move by himself
- o To prevent fatigue and injury
- o To maintain good body alignment

Practice Guideline

- Maintain functional client body alignment. (Alignment is similar whether client is standing or in bed.)
- Maintain client safety.
- Reassure the client to promote comfort and cooperation.
- Properly handle the client's body to prevent pain or injury.
- Follow proper body mechanics.
- Obtain assistance, if needed, to move heavy or immobile clients.
- Follow specific physician orders.
- Do not use special devices (e.g. splints, traction unless ordered)

Turning the Patient to a Side-lying Position

Supplies and Equipment

- Pillows
- Side rails
- Cotton blanket or towels, rolled for support

Procedure/Steps

1. Wash your hands

2. Explain the procedure to the client
3. Adjust the bed to a comfortable height
4. Lower the client's head to as flat a position as he or she can tolerate, and lower the side rail.
5. Move the client to the far side of the bed. Raise the side rail.
6. Ask the client to reach for the side rail
7. Assume a broad stance, tensing your abdominal and gluteal muscles. Roll the client toward you.
8. Position the client's legs comfortably.
 - (a) Flex his or her lower knee and hip slightly.
 - (b) Bring his or her upper leg forward and place a pillow between legs.
9. Adjust the client's arms
 - (a) Shift his or her lower shoulder toward you slightly
 - (b) Support his or her upper arm on a pillow
10. Wedge a pillow behind the client's back. Use rolled blankets or towels as needed for support.
11. Lower the bed, elevate the head of the bed as the client can tolerate, and raise the side rail.
12. Wash your hands.

Joint Mobility and Range of Motion



- a. Support the client's elbow with one hand and grasp the client's wrist with your other hand.
 - b. Raise the client's arm from the side to above the head.
 - c. Perform internal rotation by moving the client's arm across his or her chest.
 - d. Externally rotate the client's shoulder by moving the arm away from the client.
 - e. Flex and extend the client's elbow.
10. Perform all exercises on the client's wrist and fingers
- a. Flex and extend the wrist.
 - b. Abduct and adduct the wrist.
 - c. Rotate and pronate the wrist.
 - d. Flex and extend the client's fingers.
 - e. Abduct and adduct the fingers.
 - f. Rotate the thumb.
11. Exercise the client's hip and leg.
- a. Flex and extend the hip and knee while supporting the leg.
 - b. Abduct and adduct the hip by moving the client's straightened leg toward you and then back to median position.
 - c. Perform internal and external rotation of the hip joint by turning the leg inward and then outward.
12. Perform exercises on ankle and foot
- a. Dorsiflex and plantar flex the foot

- b. Abduct and adduct the toes
 - c. Evert and invert the foot
13. Move to the other side of the bed and repeat exercise.
 14. Position and cover the client. Return the bed to low position.
 15. Wash your hands.
 16. Document completion of PROM exercise.


Controlling Postural Hypotension

- o Sleep with the head of the bed elevated (8-12 inches). This makes the person's position change on rising less severe.
- o Avoid sudden changes of position. Arise from bed in three steps:
 - ⇒ Sit on the side of the bed with legs dangling for 1 minute
 - ⇒ Stand with core holding on to the edge or the bed or another non mobile object for 1 minute
 - ⇒ Sit up in the bed for one minute

Gradual change in position stimulates renin, kidney enzyme that has a role in regulating BP and which prevents a dramatic drop in BP

- o Balance is maintained with minimal effort when the base of support is enlarged in the direction in which the movement will occur

- o Contracting muscles before moving an object lessens the energy required to move it
- o The synchronized use of as many large muscles groups as possible during an activity increases overall strength and prevents muscle fatigue and injury
- o The closer the line of gravity to the center of the base of support, the greater the stability
- o The greater the friction against the surface beneath an object the greater the force required moving the object. (Pulling creates less friction than pushing)
- o



Guideline for Positioning the Client

Positioning the Client for Comfort

- Maintain functional client body alignment. (Alignment is similar whether the client is standing or in bed.)
- Maintain client safety.
- Reassure the client to promote comfort and cooperation.
- Properly handle the client's body to prevent pain or injury.
- Follow proper body mechanics.
- Obtain assistance, if needed to move heavy or immobile clients.
- Follow specific orders.
- Do not use special devices (e.g. Splints, traction) unless ordered client positioning for examination and treatment.










Figure 5 Prone position

3. Prone Position: - is used to examine the spine and back. The client lies on the abdomen with head turned to the side for comfort. The arms are held above the head or along side the body. Cover the client with a bath blanket for privacy. Caution: Unconscious clients, pregnant women, clients with abdominal incisions, and clients with breathing difficulties cannot lie in this position.




Figure 6 Sim's position

4. Sims' Position: - This position is used for rectal examination. The client rests on the left side, usually with a small pillow under the head. The right knee is flexed against the abdomen, the left knee is flexed slightly, the left arm is behind the body, and the right arm is in a comfortable position. Cover the client with a bath blanket. Caution: The client with leg injuries or arthritis often cannot assume this position




Figure 7 Fowler's position

5. Fowler's Position: - this position is used to promote drainage or to make breathing easier. Adjust the head rest to the desired height, and raise the bed section (Gatch bed) under the client's knees. Place a rolled pillow between the client's feet and use the foot of the bed as a brace, if desired. Caution: Observe for signs of dizziness or faintness when you raise the head of the bed.





Figure 8 Knee -Chest position

6. Knee-chest Position: - is used for rectal and vaginal examinations and as treatment to bring the uterus into normal position. The client is on the knees with the chest resting on the bed and the elbow rested on the bed, or with the arms above and become tense to the thighs.

7. Dorsal Lithotomy Position: - is used for examination of pelvic organs. It is similar to dorsal recumbent position, except that the client's legs are well separated and the knees are acutely flexed. The nurse will usually place the client's feet in stirrups. Keep the client covered as much as possible for privacy.

Figure 9 Lithotomy position



Crutch Walking

Crutches: - are walking aids made of wood or metal in the form of a shaft. They reach from the ground to the client's axillae.

Application of Nursing Process

Assessment

- Assess physical ability to use crutches and strength of the client's arm back, and leg muscle.
- Observe client's ability to balance self.
- Note any unilateral or unusual weakness or dizziness.
- Assess which gait is appropriate for client.
- Assess client's understanding of crutch-walking technique.

Planning/Objective

- To improve client's ability to ambulate when he/she has lower extremity injury.
- To increase muscle strength, especially in arms and legs.
- To increase feeling of well-being when client can ambulate.
- To promote joint mobility.

Implementation/Procedure

- Teaching muscle- strengthening exercises
- Measuring client for crutches

- Teaching crutch walking: Four-point gait, Three-point gait, two-point gait.
- Teaching Swing-To-Gait and Swing-Through Gait
- Teaching upstairs and downstairs ambulation with crutches.

Evaluation/Expected Outcomes

- Client's ability to ambulate is improved.
- Muscle strength of client's arms and legs is improved
- Client experiences a feeling of well-being.

Teaching Techniques of Crutch Walking

A. *Four-Point Gait*

Equipment

- Properly fitted crutches
- Regular, hard soled street shoes
- Safety belt, if needed

Procedure

1. Explain the rationale for the procedure to the client
 - a. The gait is rather slow but very stable
 - b. The gait can be performed when the client can move and bear weight on each leg.
2. Demonstrate the crutch foot sequence to the client.
 - a. Move the right crutch



Procedure

1. Explain the rationale of the procedure
 - a. The gait can be performed when the client can bear little or no weight on one leg or when the client has only one leg.
 - b. This gait is fairly rapid and requires strong appear extremities and good balance.
2. Demonstrate the crutch-foot sequence to the client.
 - a. Two crutches support the weaker extremities
 - b. Balance weight on the crutches
 - c. Move both crutches and affected leg forward
 - d. Move unaffected leg forward
3. Assess the client's progress, and correct any mistakes as they occur.
4. Remain with client until cutch safety is ensured.



Figure 11. Three-Gait Point

C. Two Point-Gait




Figure 12. Two- Gait Point

Procedure

1. Explain the procedure to the client.
 - a. This procedure is a rapid version of the four point gait
 - b. This gait requires more balance than the four gait
2. Demonstrate the crutch-foot sequence to the client.
 - a. Advance the right foot and left crutch simultaneously
 - b. Advance the left foot and the

Teaching Swing-To-Gait and Swing through Gait

Equipment

1. Properly fitted crutches
2. Regular, hard soled street shoes

Procedure

1. Explain the rationale for the procedure the client.
 - a. These gaits are usually performed when the client's lower extremities are paralyzed.
 - b. The client may use braces.
2. Demonstrate the crutch-foot sequences to the client
 - a. Move both crutches forward
 - b. Swing to gait: left and swing the body to the crutches
 - c. Swing through gait: left and swing the body past the crutches
 - d. Bring crutches informed of the body and repeat.
3. Help client practice the gait
4. Assess the client's progress and correct any mistakes as they occur.

Teaching up stairs and down stairs ambulation with crutches

Equipment

- Properly fitted crutches
- Regular, hard soled street shoes
- Safety belt, if needed

Procedure

1. Explain the rationale of the procedure to the client.
2. Apply safety belt if client is unsteady or requires support.
3. Demonstrate the procedure using a three-point gait.


Going Down Stairs

- a. Start with weight on uninjured leg and crutches on the same level.
- b. Put crutches on the first step
- c. Put weight on the crutch handles and transfer unaffected extremity to the step where crutches are placed.
- d. Repeat until the client understands the procedure

Going Upstairs

- a. Start with the crutches and unaffected extremity on the same level.

- b. Put weight on the crutch handles and lift the unaffected extremity on the first step of the stairs.
 - c. Put weight on the unaffected extremity and lift other extremity and the crutches to the step.
 - d. Repeat until client understands the procedure.
4. Help the client practice
5. Make sure that the client has adequate balance. Be ready to assist if necessary.
6. Assess the client's progress, and correct any mistakes as they occur.
7. Document the following points:



Procedure

1. Wash your hands
2. Explain the procedure to the patient
3. Position the wheelchair next to the bed or at 45° angles to the bed. Lock the wheel brakes and remove the foot rests or move them to the “up” position.
4. Prepare to move the client:
 - a. Assist the client with putting on robe and slippers.
 - b. Obtain help from another person if the client is immobile, heavy, or connected to multiple pieces of equipment.
5. Raise the head of the bed so that the client is in the sitting position.
6. Assist the client to sit on the side of the bed
 - a. Support the head and neck with one arm.
 - b. Use your other arm to move the client's leg over the side of the bed.
 - c. Allow the client's feet to rest on the floor.
 - d. Maintain the client in this position for a short-time



Study questions

- State the principle underlying proper body mechanics and relate a nursing consideration.
- State the purposes of range of motion exercise.
- Identify principles related to safe movement of clients in and out of bed.
- Demonstrate the ability to move a partially mobile client safely from bed to chair and back.
- Demonstrate the ability to teach each of the crutch walking gaits to a client.
- Mention different positions used for various examination and treatment.



Electrolyte Composition of the Fluid

Electrically charged particles act as a conductor of electrical current in the solution. E.g. $\text{NaCl} \rightarrow \text{Na}^+ + \text{Cl}^-$

Intracellular fluid and extra cellular fluid are separated by cell membrane, which is semi permeable. Body fluid composed of water, electrolyte, and non-electrolyte. The difference is maintained by the cells, which actively reject certain electrolytes, and retain others. E.g. Na^+ is reach higher in concentration in extra cellular fluid. The difference is maintained by cellular action referred as sodium pump, which reject sodium from the cells. The major ions of cellular fluid in order of their quantity are:

	<u>ICF</u>	<u>ECF</u>
K^+	141 M Eg/L	4 M. Eg/L
Mg^{++}	58 M Eg/L	2 M. Eg/L
Po4^{++}	75 M Eg/L	10 M. Eg/L
Na^+	10 M Eg/L	142 M. Eg/L
Cl^+	4 M Eg/L	103 M. Eg/L

Transport Mechanism of Electrolyte

1. Osmosis
2. Diffusion

Substances are transported between cellular and extracellular fluids between biological membranes. These transport mechanisms are mentioned above.

Osmolarity – refers to the concentration of active particles per unit of solution. Two opposing forces exist within the vascular compartment. These are:

1. Hydrostatic pressure of the blood which forces fluid out through semi permeable membrane
2. Osmotic pressure of the blood protein (colloid osmotic pressure) – which is pulling or holding force opposing the flow of fluid across the vascular membrane

When blood enters the arteriolar end of the capillaries hydrostatic pressure is greater than osmotic pressure and fluid filters out of the vessels. The movement of fluid out of the vessel is facilitated also by negative hydrostatic pressure – sucking fluid from plasma and the osmotic pressure in the interstitium.

At arterial end of capillaries, there is outward force =

$$\text{CHP} - \text{POP} + \text{Int.H.P} - \text{Int.O.P}$$

$$30 - 28 + 6 - (-5.3) = 13.3$$

At the venous end: $\text{POP} - \text{CHP} + \text{Int.H.P} - \text{Int.O.P}$

$$28 - 10 + 6 - (-5.3)$$

$$= 6.7$$

In extracellular fluid the principal osmotic forces are exerted by sodium and chlorine ions. Potassium, magnesium and phosphorous are mainly responsible for osmotic pressure within the cells.

Effect of osmosis as applied to different extracellular solute concentration will give isotonic, hyper tonic and hypotonic solution.

When all contributions to osmolality are summed the total serum osmolality ranges from 275 mosm/kg to 290 mOsm/kg.

Solutions can be categorized according to how their osmolality compared with that of extracellular fluids. When the osmolality is the same as extracellular fluid, a solution is labelled isotonic. Such a solution remains within extracellular compartment. One third is distributed to the vascular space and two thirds to the interstitial space.

A fluid with a lower or higher osmolality is labelled hypotonic or hypertonic respectively. Hypotonic fluids are distributed in proportion of to the extracellular compartment and of intracellular compartment. They are associated with cell swelling. When

hypertonic fluids are added to the vascular space, the extracellular



hydroxyl ions in a chemical reaction. The acidity or alkalinity of a solution depends upon the concentration of hydrogen ions and hydroxyl ions. A compound that completely dissociates its hydrogen ions is referred as strong acid. E.g. H_2SO_4 , HCl , H_3PO_4

A compound that partially releases its hydrogen ions is referred as weak acid. E.g. H_2CO_3 , citric acid, acetic acid.

Acid-Base Regulation

Body fluid normally have a PH of 7.35 – 7.45. The chief acid regulating from Metabolism is H_2CO_3 which is formed by a combination of CO_2



Buffers are substances, which tends to stabilize or maintain the constancy of the PH of a solution when an acid or a base is added to it.

Example: $\text{HCl} + \text{NaCO}_3 \rightarrow \text{H}_2\text{CO}_3 + \text{NaCl}$

They do this by rapidly converting a strong acid or base to a weaker one, which does not dissociate as rapidly.

Strong base $\text{NaOH} + \text{HCO}_3$ gives H_2ONaCO_3

Example of Buffering System:

- A) Bicarbonate Buffering System
- B) Phosphate Buffering System

Respiratory Regulation of Acid-Base Balance

Carbondioxide is constantly produced in cellular metabolism and diffuses from the cells into the blood and crythrocyte, and as a result CO_2 is in greater concentration in the blood. When it enters pulmonary capillaries than in the air in alveoli of the lungs.

Kidney Regulation

The kidneys play an important role in maintaining acid base balance by excretion of H^+ and forming hydrogen carbonate.

The cell of the distal tubules is sensitive to the changes in the PH.

- Excessive administration of drugs e.g ASA, Ammonium Chloride
- Renal Failure
- Dehydration
- Sever diarrhea and vomiting

Common signs and symptoms for respiratory acidosis

- Restlessness, apprehensive, slow mental response, weakness, headache, confusion " coma. PH is < 7.35
- Decreased bicarbonate
- Increased arterial CO_2 and decreased O_2
- Increased urine acidity
- Increased ammonium in urine
- Low PH in urine

Metabolic Acidosis

Headache, fatigue, drowsiness

Serum PH < 7.35

Serum bicarbonate is low

Depression in CNS

Increased respiration

Nursing Intervention

- Improve respiratory ventilation (e.g., administer bronchial dilators, antibiotic oxygen as ordered.
- Maintain adequate hydration (2 to 3 L of fluid perday)
- Carefully regulate mechanical ventilation if used.

- Monitor fluid in take and out put, vital signs, arterial blood gases (ABGs), and PH

Metabolic acidosis

Nursing interventions

- Monitor Arterial Blood Gases values
- Administer IV sodium bicarbonate carefully if ordered
- Correct underlying problem as ordered

Alkalosis: - is acid-base imbalance in which there is a decrease in H^+ concentration below 35 n mol/L and an increase in the P^H in excess of 7.45 due to carbonic acid deficit or an excess amount of bicarbonate (HCO_3).

Types of Alkalosis

1. Respiratory Alkalosis
2. Metabolic Alkalosis

Respiratory Alkalosis

Causes

1. Hyperventilation (excessive loss of carbolic acid) related to anxiety, hysteria, CNS disease which causes over stimulation of respiratory center
2. High fever
3. Hypoxia
4. Sever pain
5. High altitude


Sign and Symptoms

- Serum PH > 7.45
- Serum bicarbonate decreases
- Serum hydrogenion < 35 n mol/L
- Serum potassium decreased
- Cardiac arrhythmia
- Increased Na⁺ and K⁺ excretion in urine
- Decreased chloride ion and hydrogenion excretion
- Hyperventilation
- Increased rate and depth of respiration
- Decreased arterial blood CO₂
- Dizziness, tetany, muscle spasm (carpopedal spasm)
- Cramp, tingling in extremities
- Convulsion

Nursing Interventions

Sign and system

- Serum PH > 7.45
- Serum H+ < 35 n mal/L
- Increase serum bicarbonate
- Decreased serum potassium
- Cardiac arrhythmia
- Hypoventilation
- Slow, shallow respiration
- Increased PaCO₂ or normal
- Decreased PaO₂ if prolonged alkalosis





current philosophy is that no good foods or bad foods exist, and that



Table 1. A guideline for healthy diet

<u>Guide Line</u>	<u>Rationale</u>
– Eat a variety of foods	- No single food supplies all 40-plus essential nutrients in amounts needed variety also helps reduce the risk of nutrient toxicity and accidental contamination
– Balance the food you eat with physical activity – maintain or improve your weight	- Excess weight increases the risk of numerous chronic diseases. Such as hypertension, heart disease, and diabetes
– Choose a diet with plenty of grain products, vegetables, and fruits	- Plant foods provide fiber, complex carbohydrates, vitamins, minerals, and other substances important for good health

Therapeutic Nutrition

Therapeutic nutrition is a modifica



- To establish a means for suctioning stomach contents to prevent gastric distention, and vomiting.
- To remove laboratory contents for laboratory analysis
- To lavage (wash) the stomach in case of poisoning or overdose of medication

Equipment

- Large or small bore tube (plastic or rubber)
- Solution basin filled with warm water (if plastic tube is used) or ice (if rubber tube is used)
- Adhesive tape (2.5 cm wide)
- Disposable gloves
- Water soluble lubricants
- Facial tissues
- Glass of water and drinking straw or medication cup with water
- 20 to 50 ml syringe with an adaptor
- Basin
- Stethoscope
- Clamp (optional)
- Suction apparatus (if required)
- Gauze square or plastic specimen bag and elastic band
- Safety pin and elastic band
- Infant seat, towel, or pillow



10. Determine how far to insert
 - Use the tube to mark off the distance from the tip of the client's nose to the tip of the ear lobe and from the tip of the ear lobe to the tip of the sternum. This length approximate the distance from the nares to stomach.
 - For infants and young children, measure from the nose to the tip of the ear lobe and then to the xiphoid process.
 - Mark this length with adhesive tape, if the tube does not have marking.
11. Lubricate the tip of the tube well with water solution lubricant or water to ease insertion.
12. Insert the tube with its natural curve toward the client in to the selected nostril. Ask the client to hyper extend the neck, and gently advance the tube toward the nasopharynx. Do not hyper-extend or hyper -flex an infant neck
13. Direct the tube along the floor of the nostril and toward the ear on that side.
14. If the tube meets resistance, withdraw it, rubricate it and insert it in the other nostril. (The tube should never be forced against resistance)
15. Once the tube reaches the oropharynx (throat) the client will feel the tube in the throat and may gag or retch. Ask the client to tilt the head forward and encourage the client to drink and swallow. If the client gags, stop passing the tube momentary.

Have the client rest, take a few breaths, and take sips of water to calm the gag reflex.

16. In the cooperation with the client, pass the tube 5 to 10 cm (2 to 4 in) with each swallow, until the indicated length is inserted.
17. If the client continuous to gag and the tube does not advance with each swallow, with draw it slightly, and inspect the throat by looking through the mouth. (The tube may be coiled in the throat. If so withdraw it until it is straight, and try again to insert it).
18. As certain correct placement of the tube:

- Aspirate stomach content, and check their acidity.



- For infants or small children, tape the tube to the area between the end of the nares and the upper lip, as well as to the cheek.
20. Attach the tube to the suction source or feeding apparatus as ordered, or clamp the end of the tubing.
 21. Secure the tube to the client's gown. Loop an elastic band around the end of the tubing, and attach the elastic band to the gown with a safety pin or attach a piece of adhesive tape to the tube, and pin the tape to the gown.
 22. Document relevant information, means by which correct placement was determined and client responses.
 23. Establish a plan for providing daily nasogastric tube care
 - Inspect the nostril for discharge and irritation
 - Clean the nostril and tube with moistened cotton tipped applicators
 - Apply water-soluble lubricant to the nostril if it appears dry or encrusted.
 - Change the adhesive tape as required
 - Give frequent mouth care
 24. If suction is applied, ensure that the patency of both the nasogastric and suction tubes is maintained
 25. Document all relevant information:
 - Type of tube inserted
 - Date and time of tube insertion
 - Type of suction used

- Color and amount of gastric contents
- Client tolerance of the procedure

NASOGASTRIC TUBE FEEDING



- Feeding pump (optional)

Procedure/Intervention

1. Prepare the client and the feeding
 - Explain the patient about the feeding
 - Provide privacy
 - Position the patient in Fowler's position in bed or sitting position in a chair
 - Position a small child or infant in your lap, and provide a pacifier during feeding
2. Assess tube placement. Attach the syringe to the open end of the tube, aspirate alimentary secretions. Check the PH.
3. Assess residual feeding contents
 - Aspirate all the stomach contents, and measure the amount prior to administering the feeding. If 50 mL or more undigested formula is withdrawn in adults, or 10 ml or more in infants, check with the nurse in charge before proceeding.
 - Reinstill the gastric contents in to the stomach if this is the agency or physician's practice. Remove the syringe bulb or plunger, and pour the gastric contents via the syringe in to the nasogastric tube.

4. Administer the feeding

Before administering feeding:

- a) Check the expiration date of the feeding
- b) Warm the feeding to room temperature

Bulb syringe

- Remove the bulb from the syringe, and connect the syringe to a pinched or clamped nasogastric tube



- Instill 60 mL of water the feeding tube
 - Be sure to add the water before the feeding solution has drained from the neck of a bulb syringe or from the tubing of an administration set. Before adding water to a feeding bag or prefilled tubing set, first clamp and disconnect both feeding and administration tubes.
6. Clamp and cover the feeding tube
- Clamp the feeding tube before all of the water is instilled
 - Cover the end of the feeding tube with gauze held by an elastic band
7. Ensure client comfort and safety
- Pin the tubing to the clients gown
 - Ask the client to remain sitting upright in Fowler's position or in slightly elevated right lateral position for at least 30 minutes.
8. Dispose of equipment appropriately
- If the equipment is to be reused, wash with soap and water so that it is ready for reuse.
 - Change the equipment every 24 hours or according to the agency's policy.

9. Document all relevant information

- Document the feeding, including amount, and kind of solution taken, duration of feeding and assessment of client.
- Record the volume of the feeding and water administered on the client's intake and out put record.

10. Monitor the client for possible problems:

- Carefully assess clients receiving tube feeding for problems
- To prevent dehydration, give the client supplemental water in addition to the prescribe tube feeding as ordered.

TOTAL PARENTRAL NUTRITION

Parental nutrition is a method where by nutrients may be introduced into the system via the enteral route. It is also referee to as intravenous hyperalimenation (IVH). By passing the normal gastro intestinal system, this route provides a nitrogen source for those unable to ingest protein, carbohydrates (adequate caloric), or fats. A balanced blend of nutrients, including vitamins and minerals, can be administered peripherally, using isotonic concentrations of glucose, crystalline aminoacids, and fats; or because the solution may be irritating to the veins, nutrients can be administered through a central, high-flow vein. Hypertonic glucose, along with crystalline

aminoacids, fats, electrolytes, vitamins and trace elements is given through central vein access.

The technique requires especial handling and management of the client and the most expensive method of feeding.

It should be used only if the intestines do not work adequately, if the client has an obstruction or has fistula, if the bowel rest is required.

Implantable vascular access devices are placed under the skin in a subcutaneous pocket and a surgically tunneled silicone catheter is placed in the cephalic or external jugular vein and threaded to the superior vena cava.

Application of Nursing Process

Assessment

- Complete physical assessment and client history
- Assess weight and take a weight history
- Identify and condition that would affect TPN (renal or cardiac disease)
-

- Check label of solution with physician's orders
- Check rate of infusion on physician's order
- Check rate of infusion on physician's order
- Assess ability of client to understand instructions during the procedure.
- Ensure potency of the central venous line following the insertion
- Observe catheter insertion site for signs of infection, thrombophlebitis, or possible infiltration.
- Inspect dressing over central line to ensure a dry, non contaminated dressing.

Planning/Objective Setting


- To provide a nitrogen source for clients unable to ingest protein normally.
- To provide adequate calories for clients unable to tolerate oral feedings.
- To provide nutrients for clients requiring by pass of the gastrointestinal tract.
- To provide increased calories where regular IV solutions are insufficient.
- To prevent or correct a deficiency of essential fatty acids.
- To provide a contamination free mode of delivering the hyperalimentation solution

Implementation

- Assisting with catheter insertion
- Maintaining central vein Infusions
- Changing parenteral hyperalimentation Dressing and Tubing
- Maintaining Hyperalimentation for children

Evaluation/Expected Outcome

- Dressing remains dry and intact during interval between changes.



- Aseptone solution (optional)
- Sterile 4x4 gauze pads
- Sterile gloves
- Sterile towels or drapes
- Sterile gloves Trterile 8T20 1 Tuze pads



6. Assemble IV insertion tray or kit, normal saline solution bottle or DW, IV tubing, extension tubing, and filter.
7. Wash hands
8. Flush IV tubing with IV solution
9. Place IV tubing through infusion pump
10. Place catheter insertion equipment on bedside stand

Procedure

1. Position client in head-down position with head turned to opposite direction of catheter insertion site. Place a small roll between client's shoulders to expose insertion site.
2. Cleanse insertion area with Betadine solution (if allergic to Betadine solution, use 70% isopropyl alcohol).
3. Assist the physician to gown, put on mask and gloves prior to beginning procedure.
4. Don mask and sterile gloves.
5. Assist physician as needed during catheter insertion
6. Instruct client in Valsalva's Maneuver when stylet is removed from catheter and when IV tubing is connected to catheter
 - a) Instruct client to exhale against a closed glottis.
 - b) If client is unable to do this compress client's abdomen. Both these procedures help decrease possibility of air embolism.

7. After tubing is connected, instruct client to breath normally.
8. Tape area between tubing and catheter hub.
9. Turn on IV infusion pump, using normal saline solution, at slow rate, 10 gtt/mincle, until X-ray ensures accurate catheter placement.
10. Place Betadine over catheter insertion site. Apply transparent dressing.
11. Order portable chest X-ray to verify correct catheter placement.
12. Following confirmation of Catheter placement. Change IV solution to hyper alimentation solution and adjust flow rate as ordered.
13. Time tape the bottle after adjusting flow rate. Be prepared to document on IV hourly infusion recorded.
14. Observe for signs of complication.
15. Take vital signs every 4 hours. If signs change or temperature rises significantly, the client may be developing complicationo

- Weigh client daily – observe for fluid gain or loss. Weight gain may indicate fluid overload rather than increased nutritional gain.



UNIT FIVE

CHAPTER TWEELEVE

ELIMINATION OF GASTROINTESTINAL AND URINARY OUTPUTS

Learning Objective

At completion of the unit the learner will be able to:

- Define enema.
- List purposes of gastric aspiration, lavage, enema and catheterization.
- Mention types of enema.
- Provide enema according to its purpose and need.
- Explain mechanism of action of fluids used for enema.
- Explain purpose of catheterization.
- Identify different types of catheters.
- Describe indication of catheterization.
- Demonstrate sterility technique through out the catheterization.
- Intervene the procedure for those in need of it with understanding of both male and female catheterization.
- Identify important precautions of the procedure.


Key Terminology

anuria	dysuria	melena	projectile vomiting
constipation	enema	micturation	urgency
cystitis	fecal impaction	nocturia	urinary catheter
defecation	flatus	oliguria	urinary frequency
diarrhea	incontinence	polyuria	urinary retention
voiding	vomitus		

I Gastric Lavage

Definition- This is the irrigation or washing out of the stomach.

Purpose




- Solution as prescription/usually to care for acidic poisoning.
We use sodium bicarbonate 1 teaspoon to 500 cc. of water at a temperature of 37⁰c - 38⁰c.)
- Small jug to carry solution to the funnel
- Lubricant e.g. liquid paraffin
- Bowl for gauze swabs
- Cape or protective material to put around the patient chest
- Pail to receive returned fluid
- Mackintosh or paper to protect the floor beneath the pail
- Receiver for used esophageal tube
- Paper bag for waste material
- A tray for mouth wash after lavage
- Denature cup.
- A receiver for pt's dentures. If any, and should be labeled with the pt's name
- A receiver containing mount gag, tongue depressor, and tongue forceps if patient is unconscious
- Mackintosh to protect bed linen
- Litmus paper
- Specimen bottle. If laboratory test is required
- Measuring jug

Procedure

1. Explain procedure to the pt and ask him/her to remove artificial dentures, If any.
2. Protect pt with cape or towel


3. Protect bed linen by spreading the mackintosh on the accessible side of the bed.
4. Place mackintosh or paper under the pail to protect the floor
5. Elevate head of the bed if pt is conscious and the condition permits. But if unconscious, place in prone position with head over the edge of the bed or head lower than the body.
6. Measure the tube from the tip of the nose up to the ear lobe and from the bridge of the nose to the end of the sternum. (32



2. Gastric Lavage Using a Tube with a Bulb

Procedure

1. Clamp tubing below bulb.
2. With right hand, squeeze bulb this forcing the air out through the funnel.
3. With left hand, pinch tubing over bulb and at the same time releasing bulb. This creates a suction, which will draw the stomach contents in to the bulb.
4. Lower funnel and allow excess gastric contents to drain in to the pail.



II. GASTRIC ASPIRATION

- Aspiration is to withdrawal of fluid or gas from a cavity by suction

Purpose

1. To prevent or relieve distention following abdominal operation
2. In case of gastrointestinal obstruction, to remove the stomach or gastric contents
3. To keep the stomach empty before on emergency Abdominal operation is done
4. To aspirate the stomach contents for diagnostic purposes

There are two type of gastric Aspiration

1. Intermittent method: - In this case, Aspiration is done as condition requires and as ordered.
2. Continues method: - Attached to a drainage bag

There are 2 ways of supplying suction

- a. Simple suction by the use of a syringe
- b. An electric suction machine

- Gallipots with lubricant e.g. liquid paraffin or vase line, to lubricate the nostrils
- Gauze swabs in a bowl
- Sodium bicarbonate solution or saline to clean the nostrils
- Litmus paper
- Water in a galipot to test the right position of the tube in the stomach
- Two test tubes and laboratory forms of necessary
- Saline or plain water in a galipot to be injected, in case the stomach content is too thick to come out through the syringe.
- Rubber mackintosh and towel to protect the patient's chest.
- Receiver for soiled swabs

Procedure

1. Explain procedure to patient, in order to gain her/his co-operation
2. Prop up in an upright position with help of back rest and pillow
3. Cleanse and lubricate the nostrils
4. Lubricate the Ryle's tube with water
5. Insert the tube as directed in nasal feeding and ask the patient to swallow as the tube goes down.
6. Instruct patient to open her or his mouth to make sure the tube is in the stomach
7. After being sure that the tube is in the right position, inject about 15-20 cc. of saline or water in to the stomach.
8. Draw plunger back to with draw the fluid collect specimen, If needed

9. If the Ryle's tube is to be left in site then a spigot or clamp is used to close the end, but if it is for one aspiration and to be removed immediately, it should be withdrawn very gently to avoid irritating the mucous lining.

N.B

1. *Special care of the nose and mouth to prevent dryness should be considered*
2. *Always measure the amount withdrawn accurately noting color, contents and smell*
3. *Record on the fluid chart properly*
4. *Report any change in patient condition regarding pulse, Temperature, B.P fluid out put.*

III. Enema

Enema: is the introduction of fluid into rectum and sigmoid colon for
N.00.02 /a

Mechanisms of some solutions used in enema

1. Tap water: increase peristalsis by causing mechanical distension of the colon.
2. Normal saline solution
3. Soap solution: increases peristalsis due to irritating effect of soap to the luminal mucosa of the colon.
4. Epsom salt: The concentrated solution causes flow of ECF (extra cellular fluid) to the lumen causing mechanical distension resulting in



Guidelines

Enema for adults are usually given at 40-43°C and for children at 37.7 °C

Hot – cause injury to the bowel mucous

Cold – uncomfortable and may trigger a spasm of the sphincter muscles

The amount of solution to be administered depends on:

- Kind of enema
- The age of the person and
- The person's ability to retain the solution

Age	Amount
18 month	50-200 ml
18 mon-5 yrs	200-300 ml
5-12 yrs	300-500 ml
12 yrs and older	500-1,000 ml

The rectal tube should be appropriate: is measured in French scale

Age	Size
Infants/small child	10-12 fr
Toddler	14-16 fr
School age child	16-18 fr
Adults	22-30 fr


Purpose

- Ã To stimulate peristalsis and remove feces or flatus (for constipation)
 - Ã To soften feces and lubricate the rectum and colon
 - Ã To clean the rectum and colon in preparation for an examination. E.g. Colonoscopy
 - Ã To remove feces prior to a surgical procedure or a delivery
 - Ã For incontinent patients to keep the colon empty
 - Ã For diagnostic test
- E.g. before certain x-ray exam – barium enema
- Before giving stool specimen for certain parasites

Procedure

- Ã Inform the patient about the procedure
- Ã Put bed side screen for privacy
- Ã Attach rubber tube with enema can with nozzle and stop cock or clamp
- Ã Place the patient in the lateral position with the Rt. leg flexed, for adequate exposure of the anus (facilitates the flow of solution by gravity into the sigmoid and descending colon, which are on the side)
- Ã Fill the enema can with 1000 cc of solution for adults
- Ã Lubricate about 5 cm of the rectal tube – facilitates insertion through the sphincter and minimizes trauma
- Ã Hang the can = 45 cm from bed or 30 cm from patient on the stand

- Ã Place a piece of mackintosh under the bed
- Ã Make the tube air free by releasing the clamp and allowing the fluid to run down little to the bed pan and clamp open – prevents unnecessary distention
- Ã Lift the upper buttock to visualize the answer
- Ã Insert the tube
 - þ 7-10 cm in an adult smoothly and slowly
 - þ 5-7.5 cm in the child
 - þ 2.5-3.75 cm in an infant
- Ã Raise the solution container and open the clamp to allow fluid to flow
- Ã Administer the fluid slowly if client complains





Note

1. Most medicated retention enema must be preceded by a cleansing enema. A patient must rest for ½ hrs before giving retention enema
2. Elevate foot of bed to help patient retain enema
3. The amount of fluid is usually 150-200 cc
4. Temperature of enema fluid is 37.4 °c or at body
5. Kinds of solution used to supply body with fluid are plain H₂O, normal saline, glucose 5% sodabarbonate 2-5%
6. Olive oil 100-200 cc to be retained for 6-8 hrs is given for server constipation

Rectal Washout (Siphoning Enema)

(Colon irrigation or colonic flush)

- Also called enterolysis
- Is the process of introducing large amount of fluid into large bowel for flushing purpose and allow return or wash out fluid

Purpose

- To prepare the patient for x-ray exam and sigmoidoscopy
- To prepare the patient for rectum and color operation

Solution Used

- Normal saline
- Soda-bi-carbonate solution (to remove excess mucus)
- Tap water

- KMNO₄ sol. 1:6000 for dysentery or weak tannic acid
- Tr. Asafetida in 1:1000 to relieve distention

Procedure

- Insert the tube like the cleansing enema
- The client lies on the bed with hips close to the side of the bed (client assumes a right side lying position for siphoning)
- Open the clamp and allow to run about 1,000 cc of fluid in the bowel, then siphon back into the bucket
- Carry on the procedure until the fluid return is clear

Note:

- *The procedure should not take > 2 hrs*
- *Should be finished 1 hr before exam or x-ray – to give time for the large intestine to absorb the rest of the fluid*
- *Give cleansing enema ½ hr before the rectal wash out*
- *Allow the fluid to pass slowly*

Amount of solution

- 5-6 liters or until the wash out rectum fluid becomes clear

Passing a Flatus Tube

Purpose

- To decrease flatulence (sever abdominal distention)
- Before giving a retention enema

Procedure

- Place the patient in left. Lateral position
- Lubricate the tube about 15 cm
- Separate the rectum and insert 12-15 cm in to the rectum and tape it
- Connect the free end to extra tubing by the glass connector
- The end of the tube should reach the (tap H₂O) solution in the bowel
- The amount of air passed can be seen bubbling through the solution
(a funnel may be connected to free end of tube and placed in an antiseptic solution in bowel)
- Teach client to avoid substances that cause flatulent
- Leave the rectal tube in place for a period or no longer than 20 minute – can affect the ability to voluntarily control the sphincter if placement is prolonged
- Reinsert the rectal tube every 2-3 hrs if the distention has been unrelieved or reaccumulates – allows gas to move in the direction of the rectum.

III. Urinary Catheterization

Definition of catheterization: Is the introduction of a tube (catheter) through the urethra into the urinary bladder

- Is performed only when absolutely necessary for fear of infection and trauma



3. Determine appropriate catheter length by the clients gender
 - For adult male – 40 cm catheter
 - For adult females – 22 cm catheter
4. Select appropriate balloon size
 - 5 ml – for adults
 - 3 ml – for children

Catheterization Using a straight catheter

Purpose

- To relieve discomfort due to bladder distention
- To assess the residual urine
- To obtain a urine specimen
- To empty the bladder prior to surgery

Equipment

I. Sterile

- Kidney dish
- Galipot
- Gauze
- Towel
- Solution
- Lubricant
- Catheter

- Syringe
- Water
- Specimen bottle
- Gloves

II. Clean

- Waste receiver
- Rubber sheet
- Flash light
- Measuring jug
- Screen

Procedure

- Prepare the client and equipment for perineal wash
- Position the patient – dorsal recumbent (pillows can be used to elevate the buttocks in females).
- Drape the patient.
- Wash the perineal area with warm water and soap
- Rinse and dry the area
- Prepare the equipment
- Create a sterile field
- Drop the client with a sterile drape
- Clean the area with antiseptic solution.
- Lubricate the insertion tip of the catheter (5-7 cm in)

- Expose the urinary meatus adequately by retracting the tissue or the labia minora in an upward direction – female
- Retract the fore skin of uncircumcised mal.
- Grasp the penis firmly behind the glans and hold straighten the down ward curvature of vertical it go to the body – male
hole the catheter 5 cm from the insertion tip
- Insert the catheter into the urethral orifice
- Insert 5 cm in females and 20 cm in males or until urine comes
- Collect the urine – for specimen (about 30 ml)
- Pinch previous leakage
- Empty or drain the bladder and remove the catheter
- For adults experiencing urinary retention an order is needed on the amount to urine to be expelled

Note.

- If resistance is encountered during insertion, do not force it – forceful pressure can cause trauma. Ask the client to take deep breaths - relaxes the external sphincter (slight resistance is normal)
- **Dorsal Recumbent**

Ù The balloons are sized by the volume of fluid or air used to inflate them 5 ml – 30 ml (15 commonly) indicated with the catheter size 18 Fr – 5 ml.

- Test the catheter balloon
- Follow steps as insertion straight catheter
- Insert the catheter an additional 2.5 – 5 cm (1-2 in) beyond the point at which urine began to flow (the balloon of the catheter is located behind the opening at the insertion tip) – this ensures that the balloon is inflated inside the bladder and




Removal

- Withdraw the solution or air from the balloon using a syringe
- And remove gently

Study Questions

1. Define gastric lavage.
2. Mention indications of gastric lavage.
3. Define Enema.
4. State how the mechanism of action of soap solution enema exerts its function.
5. Mention conditions that differentiate between male and female catheterization.



UNIT SIX

CHAPTER THIRTEEN

MEDICATION ADMINISTRATION

Learning Objectives

At the end of this unit the students will be able to:

- Ù Describe various rout of drug administration.
- Ù Mention the general rules & care of administering medications.
- Ù Identify the parts and types of syringes and needles.
- Ù List the necessary equipments required for drug administration.
- Ù Mention the five rights before drug administration.
- Ù Locate the different sites of parentral drug administration.
- Ù Demonstrate essential steps of medication administration.
- Ù List precautions for medication administration.

Key Terminology

ampule	ohpthalimic	parentral
brand name	pharmacokinectics	trade name
capsule	pharmacology	transdermal
chemical name	potentiating	toxicity
dosage	prescription	transfusion
enteric coated	synergistic	vial
generic name	otic	z-track
infusion	tablet	

medication topical

Pharmacology is the study of drugs. Drugs are chemicals that alter functions of living organism.

Therapeutic agents are drugs or medications that, when introduced in to living organism, modify the physiologic functions of that organism.

Drug Metabolism


Drug metabolism in the human body is accomplished in four basic stages: absorption, transportation, biotransformation, and excretion. For a drug to be completely metabolized, it must first be given in sufficient concentration to produce desired effect on body tissues. When this “Critical drug concentration” level is achieved, body tissue change.

Route of Absorption



vascular surface and moderate pH level enhance the process of breaking down the drug.

Parental methods are the most direct, reliable, and rapid route of absorption. This method of administration includes intradermal, subcutaneous, intramuscular (IM) and intravenous (IV). The actual site of administration depends on the type of drug, its action, and the



enzymes into a less active and harmless agent that can be easily excreted. Most of this conversion occurs in the liver, although some conversion does take place in the lungs, kidney plasma and







Planning /setting objectives

- To administer medications using correct route
- To determine appropriate drug actions
- To identify when side effects or adverse reactions occur
- To accurately calculate drug dosages.

Implementation /Intervention

- Preparing for drug administration
- Creating a rapport with the patient
- Assembling necessary equipment
- Converting medication
- Calculating dosage as appropriate
- Following the five rights
- Using the unit Dose system
- Using the Narcotic control system

Evaluation /Expected out comes

- Medications are administered by correct route
- Medication action and side effects are identified
- Drug dosages are calculated accurately

Different Routes of Drug Administration

Ù Oral

Ù Topical

Ù Parenteral

- Intradermal
- Subcutaneous
- Intramuscularly


- Intravenous
- Ù Rectal
- Ù Vaginal
- Ù Inhalation

I. Oral Administration

Definition: Oral medication is drug administered by mouth

Purpose

- a. When local effects on GI tract are desired
- b. When prolonged systemic action is desired





- me a-062(sun5.43tan5.43ncen5.43s)2192 disolv5.43d in wter la-04(es)4(02650f130.431 -1.7305 5ipment 726x 20

Note

1. Remember the 5 R's
 - É Right patient
 - É Right medication
 - É Right route
 - É Right dose
 - É Right time
2. Always keep the bottle tightly closed.
3. Clean and keep the label of the bottle clear.
4. Keep medication away from light.
5. Check their expiration date.
6. Keep the rim of the bottle clean.
7. Give your undivided attention to your work while preparing and giving medications.



II. Suppository

Purpose

- To produce a laxative effect. (bowel movement),suppository is used frequently instead of enema since it is inexpensive.
- To produce local sedative in the treatment of hemorrhoids or rectal abscess.
- To produce general sedative effects when medications cannot be taken by mouth
- To check rectal bleeding


Equipment

- Suppository (as ordered)
- Gauze square
- Rectal glove or finger cot
- Toilet paper
- Receiver for soiled swabs
- Bedpan, if the treatment is in order to produce defecation.
- Screen
- Mackintosh and towel

Procedure

1. Check medication order.
2. Review client's medical record for rectal surgery/ bleeding.
3. Wash hands.
4. Prepare needed equipment and supplies.
5. Apply disposable gloves.

6. Identify client.
7. Explain procedure to client.
8. Arrange supplies at client's bedside.
9. Provide privacy.
10. Position client in Sims' position.
11. Keep client draped, except for anal area.
12. Examine external condition of client's anus. Palpate rectal walls.



25. Observe client for effects of suppository 30 minutes after administration.
26. Record medication administration.

Kinds of Suppositories Used:

1. Bisacodyl (Dulcolax) is commonly ordered for its laxative action. It stimulates the rectum and lubricates its contents. Normally 15 minutes is needed to produce bowel movement.
2. Glycerin or suppository for bringing about bowel movement. If soap suppository is used cut a splinter of soap 2-6 cm. long and wash it in hot water to smooth the rough edges before administration.
3. Bismuth - for checking diarrhea.
4. Opium, sodium barbitol etc. for sedation

III. Parenteral Drug Administration

A. Intradermal Injection

Definition: It is an injection given into the dermal layer of the skin (corneum)

Purpose

For diagnostic purpose

- a. Fine test (mantoux test)
- b. Allergic reaction

For therapeutic purpose

- c. Intradermal injection may also be given like in vaccination

Site of Injection

- The inner part of the forearm (midway between the wrist and elbow).
- Upper arm, at deltoid area for BCG vaccination

Equipment

- Tray
- Syringe & needle (sterile)
- Receiver
- Drug (to be injected)
- File
- Alcohol swab
- Marking pen
- Water in the bowl to rinse syringe and needle

Procedure

- Take equipment to the patient's side
- Explain procedure to patient
- Get hold of the arm & locate the site of injection.
- Clean the skin with swab and inject the drug about 0.1. 0.2 inch in to the epidermis after the bevel of the needle is no longer visible. Don't massage the site.
- Check for the immediate reaction of the skin (10-15 minutes later for tetanus, 20-30 minutes later for penicillin)
- If it is for tine test, mark the area
- Chart the data and time of the administration of the drug.
- Take care of the equipment & return to their places.

- Do not forget to do the reading after 72 hours if it is for fine test (tuberculin test)
- Document about the procedure

B. Sub - Cutaneous Injection

Definition: Injecting of drug under the skin in the sub- cutaneous tissue, (under the dermis)

Purpose:

- To obtain quicker absorption than oral administration
- When it is impossible to give medication orally

Equipment

- Tray
- Sterile syringe & needle (disposable)
- Alcohol swabs
- Medication
- File
- Medication card and patient chart
- Receiver
- Water in a bowel
- Disposing box

Site of Injection

- Outer part r.a abrm
-

Procedure

- Take equipment to the pt's bed side or room
- Explain the procedure to the patient
- Draw your medication
- Expel the air from the syringe
- Clean the site (usually it is in upper arms, thighs or abdomen)
- Grasp the area between your thumb & forefinger to tense it.
- Insert the needle elevate about 45° - 60° angle.
- Pierce the skin quickly & advance the needle
- Aspirate to determine that the needle has not entered a blood vessel
- Inject the drug slowly.
- After injecting withdraw the needle and massage the area with alcohol swab.
- Chart the amount and time of administration
-

C. Intra- Muscular Injection


Definition: It is an introduction of a drug into a body's system via the muscles.

Purpose

- To obtain quick action next to the intra- venous route
- To avoid an irritation from the drug if given through other route.

Equipment

- Tray
-








Figure 10IM injection sites A : Vastus lateralis, B. Deltoid muscle, C. Gluteal Maxmus

Procedure

- Do the ABC of the procedure.
- Prepare tray & take it to the patient's room

- Prepare the medication
- Draw the medicine
- Expel the air from the syringe
- Choose the site of injection (the site for intra- muscular)
- Using the iliac crest as the upper boundary divided the buttock into four. Clean the upper outer quadrant with alcohol swab:
- Stretch the skin and inject the medicine
- Draw back the piston (plunger) to check whether or not you are in the blood vessel (if blood returns, withdraw and get a new needle & reinject in a different spot)
- Push the drug slowly into the muscle
- When completed, withdraw the needle and massage the area with swab gently to and absorption.
- Place the patient comfortably
- Take care of the equipment you have used & return to their places
- Chart the amount, time route and type of the medicine
- Check the patient's reaction

Note:

D. I.V. INJECTIONS

Definition: It is the introduction of a drug in solution form into a vein. Often the amount

is not more than 10.ml. at a time.

Sites for IV injection

1. Dorsal Venous network
2. Dorsal metacarpal Veins
3. Cephalic Veins
4. Radial vein
5. Ulnar vein
6. Baslic vein
7. Median cubital vein
8. Greater saphenous vein

Purpose

- When the given drug is irritating to the body tissue if given through other routes.
- When quick action is desired.
- When it is particularly desirable to eliminate the variability of absorption.
- When blood drawing is needed (exsanguinations)

Equipment

- Tray
- Towel and rubber sheet
- Sterile needle and syringes in a sterile container

- Sterile forceps in a sterile container
- Alcohol swabs
- File
- Medication
- Tourniquet
- Receivers (2)
- Treatment Chart
- Glove

Procedure

- Prepare your tray & the medication
- Explain the procedure to the patient
- Position the patient properly
- Place rubber and towel under his arm(to protect the bed linen)
- Expose the arm and apply tourniquet
- Ask pt. To open and close his fist.
- Palpate the vein and clean with alcohol swab the site of the injection (Which is mainly the mid cubital vein of the arm)
- Clean with a circular motion; proceed from center of the site outward.
- Hold the needle at about 45° angles in line with the veins.
- Puncture the vein and draw back to check whether you are in the vein or not.

(Blood return should be seen if you are in the vein)

-



Purpose

- To maintain fluid & electrolyte balance
- To introduce medication particularly antibiotics.

Equipment

- IV fluid as ordered
- Sterile syringe & needle
- Rubber & towel
- Receiver
- Alcohol swabs
- Arm board
- Bandage & scissors
- Tourniquet
- I.V pole
- Adhesive tape
- Medication chart



E.g. if 1000ml of 5% D/w is to run for 24 hrs, how many drops per minutes should it run?

$$\frac{1000 \text{ ml.} \times 15 \text{ gtt/ml.}}{24 \times 60 \text{ min.}} = \frac{1000 \times 15 \text{ gtt.}}{24 \times 60 \text{ min.}} = 10 \text{ gtt/min}$$

Note:

1. The arm board should be long enough to extend beyond the wrist and elbow joint.
2. Board should be padded
3. Infusion bottle should be labeled with the date, time infusion is started, drops per minute, and any added medications. If more than one bottle as used in 24 hrs, it should be labeled as bag 1,2,3, and so on.
4. Extend the arm in the most comfortable position.
5. Usual areas used for intravenous infusion are:
 - a) The median basilic vein on the inner surface of the arm.
 - b) A vein on top of the foot
 - c) In an infant the jugular vein and the scalp vein

F. Blood Transfusion

Definition: It is the giving of blood to a patient through a vein

Purpose

- To counteract severe hemorrhage and replace the blood loss.
- To prevent circulatory failure in operation where blood loss is considerable, such as in rectal resection hysterectomy and arterial surgery.



Procedure

- Explain procedure to patient
- Before blood transfusion is administered, the nurse has to check the blood group & RH- factor if cross match of the donor's & the recipient's blood is done and is compatible. And also check for HIV other blood born pathoges.
- Prepare the tray with necessary items
- Before taking it to the patient's room, check the patient's name, hospital number, bed number, blood group, Rh. Factor and the expiry date with a 2nd nurse or a doctor.
- Blood should be used within 21 days of its withdrawal date, if sodium citrate is used it can be used until 36 days.
- Take it to the pt's room
- Hang the bottle & remove the air from the tubing
- Put pt. in a comfortable position.
- Place rubber & towel under the arm
- Check the vital signs before administering
- Choose the vein
- Apply tourniquet
- Clean the skin & feel for a distended vein & clean again.
- Puncture the vein with the needle (the needle here should be short and wide so that it does not cause occlusion easily)
- After you make sure that you are in the vein release tourniquet & open the clamp.
- The drop/minute at the beginning should be very slow
- Watch patient closely for any reaction

- If there is no reaction from the patient regulate the rate of flow according to the patient's conditions & the order.
- Splint the arm & position it comfortably.
- Remove the equipment you have used, wash and return to its proper place.
- Record the time you started the blood & any other pertinent information.
- Check pt. frequently.

Note:

1. Always member to have anti- histamine injection ready in case a patient has reaction from the blood.
2. Be familiar with the most usual symptoms of blood reactions which are:-

Immediate Reaction:

- a) Headache
- b) Backache
- c) Chills
- d) Pyrexia
- e) Rash of the skin (urticaria)

Late Reaction

- a) Dyspnea
- b) Renal shut down in severe cases
- c) Heamaturia
- d) Chest pain

G. Cut Down

Definition - Dissection of a vein for inserting I.V cannula or needle.

Purpose

- When vein puncture is difficult
- When prolonged, continuous infusion is needed
- When rapid infusion is important and emergency situation combine these indications.

Equipment

Sterile

- Dressing forceps (1)
- Cotton balls in a gallpot
- Solution for cleansing
- Gloves
- Hole sheet (Fenestrated towel)
- Syringe and needle
- Scalpel (surgical knife)
- Mosquito forceps (3)
- Aneurysm needle (1)
- Silk
- Intravenous cannula or vein flow (2)
- Small, straight scissors (1)
- Small, curved scissors (1)
- Needle holder (1)
- Round needle (1)



Administering Vaginal Medications

Purpose

- To treat or prevent infection
- To remove an offensive or irritating discharge
- To reduce inflammation
- To relieve vaginal discomfort

Equipment

- Prescribed vaginal suppository8()Tj/TT7 1 Tf-0.4611 -1.7844 TD0 Tw<0078>Tj/TT2 1 Tf0.4611
-



- A. Take suppository from wrapper and lubricate smooth or rounded end.
 - B. Lubricate gloved finger of dominant hand.
- Offer client perineal pad.

15. Apply cream or foam:

- A. Fill applicator as directed.
- B. Retract client's labial folds with nondominant gloved hand.
- C. With dominant gloved hand, insert applicator 5 to 7.5 cm; push plunger.
- D. Withdraw applicator and place it on paper towel. Wipe away lubricant from client's orifice and labia.
- E. Wash applicator and store for future use.

16. Remove and discard gloves.

17. Wash hands.

18. Instruct client to remain flat on her back for at least 10 minutes.

20. Inspect condition of client's vaginal canal and external genitalia between applications.

21. Record medication administration.

C. Retract client's labial folds with nondominant gloved hand. D. Insert rounded end of suppository 7.5 to 10 cm along posterior wall of vaginal canal.

E. Withdraw finger and wipe away lubricant from client's orifice and labia.



damp washcloth or cotton ball over eye for a few minutes.

14. Hold cotton ball or clean tissue in nondominant hand on client's cheekbone just below lower eyelid.
15. With tissue or cotton ball resting below lower lid, gently press downward with thumb or for 5 seconds. a eNrw



B. Holding ointment applicator above lower lid margin, apply thin stream of ointment evenly along inner edge of lower eyelid on conjunctiva from inner canthus to outer canthus.

C. Have client close eye and rub lid gently in circular motion with cotton ball, if rubbing is not contraindicated.

19. Intraocular disk procedures:

A. Application:

- (1) Wash hands.
- (2) Put on gloves.
- (3) Open package containing disk. Gently press fingertip against disk so it adheres to finger. Position convex side of disk on fingertip.
- (4) With other hand, gently pull client's lower eyelid away from the eye. Ask client to look up.
- (5) Place disk in the conjunctival sac so that it floats on the sclera between the iris and lower eyelid.
- (6) Pull client's lower eyelid out and over disk.

B. Removal:

- (1) Wash hands.
- (2) Put on gloves.
- (3) Explain procedure to client.
- (4) Gently pull on client's lower eyelid to expose disk.
- (5) Using forefinger and thumb of opposite hand, pinch disk and lift it out of client's eye.

20. If excess medication is on eyelid, gently wipe eyelid from inner to outer canthus.

21. If client had an eye patch, apply clean patch by placing it over affected eye so entire eye is covered.
Tape securely without applying pressure to eye.
22. Remove gloves.
23. Dispose of soiled supplies in proper receptacle.
24. Wash hands.
25. Note client's response to instillation. Ask if any discomfort was felt.
26. Observe client's response to medication by assessing visual changes and noting any side effects.
27. Ask client to discuss drug's purpose, action, side effect, and technique of administration.
28. Have client demonstrate self-administration of next dose.
29. Record drug administration and appearance of client's eye.
30. Record and report any undesirable side effects.

Administering Ear Medications

Purpose:

- To relieve pain
- To treat infection
- To better visualize during examination

Equipment

- Disposable tissues
- Medication
- Cotton ball
- Gloves

Procedure/Steps


1. Check the medication order against the original physician's order.
2. Wash hands carefully.
3. Prepare the medication following the "five rights."
4. Proceed to the client's bed side and identify the client.
5. Put on gloves
6. Ask the client to lie on the side of unaffected ear.
7. Remove excess drainage with a dry wipe.
8. Expose the external ear canal by properly adjusting the client's ear lobe. For adults, pull the lobe up, back, and outward. For children, pull the lobe down and back.

11. If the procedure is ordered for both ears, allow 5-10 minutes



The venture mask gives a controlled amount of O_2 i.e. it is not high to cause respiratory depression & it is sufficient to relieve anoxia. It gives 24-35% of O_2

The B.L.B mask provides an oxygen concentration of 90% with the flow meter set at 7 liters/minute. This kind of mask allows the patient to eat, drink and to expectorate. If the patient cannot breath through his nose, the B.L.B mask should not be used.



that the tubing is secured to the bed linen by means of safety pin. Stay with the patient till he is reassured if it is his first time to be on oxygen therapy.

2. Giving oxygen by nasal catheter.

There are different kinds of catheters,

- a) A fine catheter
- b) A spectacle frame, which carries two, places of rubber tubing and is worn by the pt.
- c) Two soft rubber catheters connected by **y- shaped** connection to the tube on O₂ apparatus.

Equipment

- Oxygen cylinder with regulating valve and pressure tubing
- Wolf's bottle
- Glass connection
- Fine catheters, lubricant, plaster
- Safety pin
- Tray containing a galipot of saline or water. Receiver for soiled applicators.

Procedure

1. Procedure is the same as giving oxygen by mask:
(Procedure 1-4)
2. Connect the fine catheter with the pressure tubing. Turn on the fine adjustment to the required rate of flow the maximum liter flow being 6-7 litter /minute.

5. No smoking sign for the unit

Procedure

1. Remove all electrical appliances from the room as this may produce sparks.
2. Post sign of no smoking on many places in the unit
3. Prepare and check if the applicator is working properly.
4. Bring the oxygen unit to the bedside and fix the tent on the hanger.
5. Close all appliances of the tent: place ice if the apparatus is without refrigeration device.
6. Tuck the side of the hold of tent under the mattress as far as they will go.
7. Fill the tent with 12-15 liters of oxygen 40-60% concentration for the first half hour.
8. After the first half hour regulate the flow of oxygen to 6-10 liters or as ordered by the doctor until the treatment is

clothing, electric pads, bells mechanical toys should be forbidden.

2. Alcohol must not be applied to the pt's skin
 3. The catheter tip and the cylinder itself must not be lubricated with Vaseline or oil or any kind
 6. Cylinders must be handled carefully as the oxygen is under pressure.
 7. The fine adjustment should always be closed when the main tap is turned on.
 8. Check that there is no obstacle in the pt's airway before firing oxygen in order to prevent pt. From suffocation.
 9. A rate of 2-liters/ minute is commonly used when oxygen used in case of emergency instead of free air. In the case of asphyxia liter/min may be needed.
- Protect patient from asphyxia, inspecting regularly pressure gauge and flow meter and noting pulse, respiration, color, mental state and necrosis from carbon dioxide.

ii. Steam Inhalation

Definition: It is the intake of steam alone or with medication through the nose or mouth

Purpose

1. In order to produce a local effect on the upper respiratory passage during cold, sinusitis, laryngitis, bronchitis etc. common drugs used are frier balsam (tincture of benzoin compound, eucalyptus. Menthol, camphor)





- Aminophylline - Tongue depressor



4. Which one of the following site of injection most preferred for young children?
 - a. Vastus lateralis
 - b. Ventrogluteal
 - c. Deltoid muscle
 - d. Dorsogluteal
5. Explain the difference between intravenous injection and intravenous infusion.
6. List at least three immediate complications of blood transfusion.
7. Define inhalation




UNIT SEVEN

CHAPTER FOURTEEN

WOUND CARE

Learning Objectives

- Differentiate types of wounds.
- Explain the purpose of wound care.
-





Purpose

- To keep wound clean
- To prevent the wound from injury and contamination
- To keep in position drugs applied locally
- To keep edges of the wound together by immobilization
- To apply pressure

Equipment

- Pick up forceps in a container
- Sterile bowl or kidney dish
- Sterile cotton balls
- Sterile galipot
- Sterile gauze
- Three sterile forceps
- Rubber sheet with its cover
- Antiseptic solution as ordered
- Adhesive tape or bandages
- Scissors
- Ointment or other types of drugs as needed
- Receiver
- Spatula if needed
- Benzene or ether.

Technique

Aseptic technique to prevent infection





- Probe and director if required
- Scissors
- Benzene or ether
- Bandages or adhesive tape
- Bucket to put in soiled dressing

Procedure

Explain procedure to the patient

- Clean trolley or tray and assemble sterile equipment on one side and surgically clean items on the other side. Make sure the tray or trolley is covered.
- Drape patient and position comfortably.
- Place rubber sheet and its cover under the patient and place



- Dress the wound and make sure that the wound is covered completely
- Fix dressing in place with adhesive tape or bandages
- Leave patient comfortable and tidy
- Cleanse and return equipment to its proper places
- Discard soiled dressings properly to prevent cross infection in the ward.

N.B.

- If sterile forceps are not available, use sterile gloves
- Immerse used forceps, scissors and other instrument in strong antiseptic solution before cleansing and discard soiled dressing properly.
- In a big ward it is best to give priorities to clean wounds and then to septic wounds, when changing dressings, as this might lessen the risk of cross infection.
- Consideration should be given to provide privacy for the patient while dressing the wound.

Equipment

- Sterile kidney dish
- Sterile gallipot
- Sterile Scissors
- 3 Sterile forceps
- Sterile cotton balls
- Sterile gauze
- Anti septic solution as ordered
- Sterile safety pins if needed
- Cotton wool or absorbent
- Receiver
- Rubber sheet and its cover
- Adhesive tape or bandage
- Dressing scissors
- Ointment paste or paraffin gauze
- Spatulas if needed
- One pair sterile gloves if available.

Procedure

Explain procedure to the patient

- Cleanse tray or trolley and organize the needed equipment and make sure it is covered.
- Drape and position the patient according to the need and put rubber sheet and its cover under the part to be dressed
- Remove the outer layer of the dressing

- Use sterile forceps and remove the inner layer of the dressing (pay attention so that the drainage tube is not pulled out with the old dressing)
- Observe the wound for the type and amount of discharge
- Clean the wound with cotton balls soaked in antiseptic solution.
- Grasp the top of the drainage tube with sterile forceps. Pull it up a short distance while using gentle rotation and cut off the tip of the drain with sterile scissors (the length to be cut depends on the instruction or order).
- Place sterile safety pin through the drainage tube close to the wound using sterile gloves or sterile gauze, if it is in the abdomen to stop the drainage tube slipping down out of sight.
- Make sure the wound and the skin around are properly cleaned.
- Apply ointment or paste to the skin with spatula directly around to prevent irritation and excoriation (if the excoriation exists use paraffin gauze to prevent further complications).
- Cut the gauze towards its center to fit around rubber drainage tube, so that it fits properly around the tube thus preventing discomfort.
- Use adhesive tape or bandages to secure the dressing in place.
- Record state of wound and the drainage.

Note.

- Safe method should be used for disposing old dressing.



- Receiver for soiled dressings

Procedure

Explain the procedure to the patient and organize the needed items.

- Drape and position patient
- Put rubber sheet and its cover under the part to be irrigated
- Remove the outer layer of the dressing
- Remove the inner layer of the dressing using the first sterile forceps.
- Put the receiver under patient to receive the out flow
- Use syringe with desired amount of solution fitted with the catheter.
- Use forceps to direct the catheter into the wound.
- First inject the solution such as H_2O_2 at body temperature gently and wait for the flow. This must be followed by normal saline for rinsing.
- Make sure the wound is cleaned and dried properly.
- Dress the wound and check if it is covered completely
- Secure dressing in place with adhesive tape or bandage
- Leave patient comfortable and tidy
- Record the state of the wound
- Clean and return equipment to its proper place.

Note:

- Keep patient in a convenient position. According to the need so

- Use sterile technique and warm solution for irrigating the wound.

Suturing

Definition: The application of stitch on body tissues with the surgical needle & thread.

Purpose

- To approximate wound edges until healing occurs
- To speed up healing of wound
- To minimize the chance of infection
- For esthetic purpose

Equipment

- Tray or trolley covered with a sterile towel
- Sterile needle holder
- Sterile round needle (2)
- Sterile cutting needle (2)
- Sterile silk
- Sterile cat- gut
- Sterile tissue forceps
- Sterile suture scissors
- Sterile cotton swabs in a galipots
- Sterile solution for cleaning
- Sterile dressing forceps
- Sterile receiver
- Sterile gauze

- Sterile plaster
- Dressing scissors
- Local anesthesia
- Sterile needle & syringes
- Sterile gloves
- Sterile hole- towel (Fenestrated towel)

Procedure

- Explain procedure to patient
- Adjust light
- Wash your hands
- Clean the wound thoroughly
- Wash your hands again
- Put on sterile gloves
- Drape the Wound with the hold- sheet
- Infiltrate the edges of the wound to be sutured with local anesthesia.
- Approximate the edges of the fascia with the help of the tissue forceps and using the round needle and cat- gut. Suture the fascia layer first.
- Using the cutting needle and silk, suture the outer layer of skin approximating the edges with the help of the tissue forceps.
- Clean with iodine and cover with sterile gauze.
- Remove the hole- Sheet
- Make patient comfortable
- Remove all equipment, wash & return to its proper place or send for sterilization.

Note:

- Do not suture wounds that are over 12 hrs old. However, such wounds have to be seen by a doctor since excision of all dead & devitalized tissue and eventual suturing may be required.
- Check that the patient gets his order for T.A.T before he leaves the hospital.
- Do not suture deep wound.
- Before you suture any wound, make sure it is free of any foreign bodies.

Removal of the Stitch

Technique: Use aseptic technique

Principles

- Sutures may be removed all at a time or may be removed alternatively.
- Do not cut stitches in more than one place as a part of it may be left behind and may cause infection.
- Suture is lifted slightly by the knot to allow scissors to go under and one part of the suturing from the cleanest part of the wound to the most contaminated part.
- Cleanse the skin around with antiseptic. Remove – gum with benzene or ether and discard the forceps
- Place sterile gauze to receive stitches.
- Take a pair of scissors in the right hand.
- Take a dissecting forceps in the left hand.
- Pull-up gently the knot resting against the skin with the forceps, pass the point of the scissors under the knot then cut the stitch on one side and remove.



Clips

Definition: Metal suture used to stitch the skin

Purpose

Same as *suturing with stitch*

Equipment

- Michel clip applier
- Clip
- Tissue forceps (toothed dissecting forceps)
- Cleaning material- same as *suturing with stitch*.

Procedure

The first part of procedure is the same as for *suturing with stitch* Except that instead of suturing the skin with thread and needle you would apply clips with the applier.

Removal of Clips

Technique

Use aseptic technique

Equipment

- Sterile gauze
- Sterile cotton balls
- Sterile kidney dish
- Sterile forceps 3

- Sterile clip removal forceps
- Antiseptic solution (Savalon 1% and iodine)
- Receiver
- Benzene or ether
- Adhesive tape or bandage

Procedure

Explain procedure to the patient and organize the needed equipment

- Drape and position patient
- Protect bedding with rubber sheet and its cover
- Remove old dressing and discard.
- Cleans wound with antiseptic solution starting from the cleanest part of the wound to the most contaminated part and discard the cotton ball.
- Place sterile gauze to receive removed clips.
- Take clip remove with the right hand and dissecting forceps with the left hand.
- Insert the lower blade of the clip remove below the middle of the clip using the dissecting forceps as a support of old the clips in place, and close the blade firmly as this will cause disagreement of the clips from the skin.
- Receive clips on sterile gauze
- Apply iodine on the skin punctures if required
- Dress the area if required
- Secure dressing in place with adhesive tape
- Leave patient comfortable and tidy
- Record the state of scare



UNIT EIGHT

CHAPTER FIFTEEN

PERI OPERATIVE NURSING CARE (PRE & POSTOPERATIVE NURSING CARE)

Learning objectives:

- Ã List steps in pre operative preparation.
- Ã Identify the high-risk surgical patients.
- Ã Describe the major assessment skills, needed in the pre operative, intra operative, and postoperative stages.
- Ã Explain the purpose of informed consent.
- Ã Perform general postoperative measures such as: obtaining vital sings, assessing level of consciousness, assessing surgical pain.
- Ã Report and document post operative complication.
- Ã Assess for patient air way.

Key terminology

anaesthesia	hypothermia	postoperative
atelectasis	hypoxia	preoperative
elective	intraoperative	suture
embolus	perioperative	
evisceration	pneumonia	



- Powerlessness

Planning/Objectives

Prior to surgery, the client:

- Demonstrates physical preparedness for surgery (absence of significance deviations from normal in vital signs, no signs of infection).
- Verbalize any concerns or fears related to the surgery.
- Provides informed consent for the surgery.
- Correctly demonstrates how to turn, deep breath, use equipment.
- Verbalizes understanding of post operative pain management program.
- Verbalizes understanding of post operative activity plan.
- Demonstrate the present of adequate caregivers at home after discharge.

Implementation

- Establish a supportive and trusting nurse-client relationship.
- Develop and implement a teaching plan that:
 - g Familiarizes the client and family with what to expect on the day of surgery.
 - g Prepares the client to participate in the pain management program.

Pre-operative

Purpose

- To prepare the patient emotionally, mentally and physically for surgery.
- To prevent any complication before, during and after surgery.


Equipment



- Be sure the patient's hair is clean. If the surgery is on the face, neck, shoulders or upper chest, the hair should be thoroughly washed, combed and tied up to keep it from touching the operative area. If the surgery is on the head the area must be shaved and the hair washed.
- If an enema has been ordered, give the night before surgery. Be sure this is given and is effective. Chart the results.

Psychological preparation

- If the patient does not yet understand what will be done. Explain briefly what the operation is and how it will help him. Avoid telling him anything that would make him worry.



afternoon., fluids and food should not be taken in the morning depending on the orders

- Check the cleanliness of body areas, umbilicus, nails and hair.
- Shave the hair from the skin of the operative area thoroughly. Some one should check to see if all the hair has been removed. Wash the skin well with soap and water before and after shaving.
- Check the orders for preoperative treatment, such as enema, catheterization of Foley catheter.
- The patient's temperature, pulse, respirations and blood pressure should be taken and recorded on the chart just before surgery.
- Give the premeditation as ordered, being careful to give the right amount at the right time to the right patient and record.

Just before surgery

- Have the patient void, if he is unable to void inform the doctor.
- Assist patient to move to the stretcher. The patient may be very sleepy or dizzy from the preoperative medications and may hurt himself. Support the stretcher to keep it from rolling as the patient moves onto it.
- Make sure his elbows are close to his sides or over his chest
-

Shaving

Purpose

- To minimize the danger of infection by decreasing the number of bacteria on the skin.



- When all the hair has been shaved off, rinse the skin with clear water. (If hair is long it could be shortened before shaving)
- Wash it again with soap, and water. Use enough soap to make lather.
- Be sure to



Face Operation

- Shave the site of the operation will be
- If the patient is a man, make sure that the face is completely free from beard.
- Wash face
- Be careful not to get soap into the patient's eyes.

Anterior Neck Operations:

- Wash the patient's head and neck
- If the patient is a woman, tie her hair, and keep it away from her neck, or cut it short.
- Shave the front and sides of the neck from the chin to the end of the sternum, and out to the shoulders.
- The area must be clean.

Posterior Neck Operations:

- The head and neck should be washed. Cut the hair short or tie out of the way.
- Shave at least 15cm. and around the place of incision.

Spinal-Operations

- Ask the doctor where the site of operation will be
- Shave at least 15-25cm. all around the area of incision.

Breast Operations

- Shave the anterior and posterior chest from neck to the waist line on the side where the surgery will be

- Shave the axilla on that side and the arm as far down as the elbow.

Kidney Operations

- Turn the patient on his side with the operative side upward.
- Shave from the sternum to the groin and across the side the same width up to the spinal column.
- Shave the axilla on that operation is a long one, so a large area must be prepared.

Abdominal operations

- Shave the whole abdomen from the end of the sternum down to the pubes.
- The umbilicus must be clean

Perennial and Rectal Operations

- The pubic and perinea hair must be shaved from the pubes to the anal area.
- Shave at least 15 cm. down the inside of the thighs both sides

Limb Operations

- The whole limb should be washed well
- Shaved at least 15 cm all around the operative area.
- If the operation is on the upper arm or the upper leg. The axilla or perineum should be shaved as well.
- If the operation is near the hand or foot cut the nails very short and clean them well.

Intraoperative Nursing Care

Observing a client undergoing surgery may be a component of a



- To rehabilitate the patient.

Equipment

- Anesthetic bed
- Oxygen
- Sphygmomanometer
- Stethoscope
- Suction machine (as needed)
- Extra rubber sheet (as needed)
- I.V stand
- Emergency drugs (to be ready in wards)
- Bed blocks (as needed) for shock

Procedure

- Prepare anesthetic bed (see section on bed making)
- Assist operating room nurse in placing patient in bed. An unconscious patient may be placed on either his right or left side unless specified
- Check post- operative orders and adjust flow of drip of IV fluid.
- Take blood pressure, pulse and respiration as ordered (usually every 15 minutes until stable)
- Encourage patient cough and breathe deeply every 15 minutes for two hours, and then every two hours until able to be up, unless other orders are written.
- Check dressing for any excessive bleeding or drainage.
- Check for tubes to be connected to drainage bottle- no kinks in tubing. Secure tubing with bedding.



- If gastric suction is present make sure it is working properly
- Frequent mouth care for patients who are not allowed to drink.

Eye Surgery

- Must lie very still because the incision and sutures can be damaged by pulling on the eye muscles. Both eyes may be covered.
- Room may be quiet and dark
- Patient must be fed.

Spinal Surgery

- Must lie on abdomen of back with bed flat, and supported by fracture board mattress.
- Patient may be in a body cast. Care must be given to prevent bed sores where the cast rubs.

Thyroidectomy

- Place in high lowers position. This will make it easier to breathe since the pressure of dressing and swelling may give choking feeling.
- An emergency tracheotomy set should always be at the bedside or nurses office for first three days, in case of hemorrhage or swelling trachea.
- The complication “Thyroid crisis” must be reported immediately as death can occur if condition is not treated quickly.

Tonsillectomy

Child

- Lie on abdomen or side to prevent blood drainage into throat, lunge or stomach.
- Watch carefully for excessive bleeding.

Adult

- If conscious, he may sit in semi- fowler's positron in order to spit the blood more easily.
- Watch carefully for excessive bleeding.

Study Questions

1. Mention the purposes of preoperative nursing care.
2. Why shaving is indicated before surgery?
3. Differentiate between the roles of scrubbed and circulating nurses in operating room.
4. State purpose of postoperative nursing care.
5. List some important equipment to provide care for immediate postoperative patients.
5. Why informed consent is required before surgery?

UNIT NINE
CHAPTER SIXTEEN
CARE OF THE DYING AND
POSTMOTREMCARE



total absence of brain activities as assessed and pronounced by the physician.

Spirituality and Death

Death often forces people to consider profaned questions: the meaning of life, the existence of the soul, and the possibility of an after life. Individuals faced with death, their close friends, and family often relies on a spiritual foundation to help them meet these challenging concepts. Spirituality takes several forms. Bernard and Schneider mention three levels of spiritual support for dying persons.

- o The first level is drawing strength from God.
- o The second level is strength generated by prayer.
- o The third level is strength from caring relationships with others.

For those whose spirituality does not include beliefs rooted in organized religion, support may take the form of compassionate care and the acceptance of personal beliefs.

Consider the spiritual dimension of your client's needs. Meeting basic human needs is an expression of caring that dying individuals will appreciate even if they can no longer communicate with you verbally.

move into depression. Dr. Ross writes about two kinds of depression. One is preparatory depression; this is a tool for dealing with the impending loss. The second type is reactive depression. In this form of depression, the person is reacting against the impending loss of life and grieves for him or herself.

The final stage of dying is that of *acceptance*. This occurs when the person has worked through the previous stages and accepts his or her own inevitable death. With full acceptance of impending death comes the preparation for it; however, even with acceptance, hope is still present and needs to be supported realistically.

Many factors influence how individuals accept death. Personal values and beliefs about life; views of personal successes, both financial and emotional; the way they look physically when experiencing the dying process; their family and friends and their families' attitudes and reactions; their past experiences in coping with difficult or traumatic situations; and, finally, the health care staff who are caring for them during this process – all affect an individual's attitude toward dying.

Nursing Process

Assessment


Observe the physical symptoms.

- Evidence of circulatory collapse
- Variations in blood pressure and pulse
- Disequilibrium of body mechanisms

- Deterioration of physical and mental capabilities
- Absence of corneal reflex

Observe the client's ability to fulfill basic needs without complete assistance.

- Assess the nature and degree of pain the client is experiencing.
- Observe for impending crisis or emergency situation.
- Observe for psychosocial condition.
- Need to establish a relationship for support
- Grief pattern and stage of grief the client is experiencing
-



2. Recognize the symptoms of urgency or emergency conditions and seek immediate assistance.
3. Notify the charge nurse if there is an impending crisis and perform emergency actions until help arrives.
4. Encourage dying clients to do as much as they can for themselves so that they do not just give up-a state that only reinforces low self-esteem.
5. Provide emotional nursing care for the client.
 - a. Form a relationship with the dying client. Be willing to be involved, to care, and to be committed to caring for a dying client.
 - b. Allocate time to spend with the client so that not only physical care is administered.
 - c. Recognize the grief pattern and support the client as he or she moves through it.
 - d. Recognize that your physical presence is comforting by staying physically close to the client if he or she is frightened. Use touch if appropriate and nonverbal communication.
 - e. Respect the client's need for privacy and withdraw if the client has a need to be alone or to disengage from personal relationships.
 - f. Be tuned into client's cues that he or she wants to talk and express feelings, cry, or even intellectually discuss the dying process.

g. Accept the client at the level on which he or she is



Care After Death



- Wash hands and wear clean gloves according to agency policy
- Close doors of the room or pull curtain
- Raise bed to comfortable working level (when necessary)
- Arrange for privacy and prevent other patients from seeing in to room.
- Close patient's eyes and nose if necessary
- Remove N.G. tubes and other devices from patient's body
- Place patient in supine position
- Replace soiled dressing with clean ones when possible
- Bath patients as necessary
- Brush or comb hair
- Apply clean gown
- Care for valuable and personal belongings and document dispersement
- Allow family to view patient and remain in room
- Attach special level if patient had contagious disease
- Await arrival of ambulance or transfer to morgue
- Remove gloves and wash hands
- Document the procedure

Study Questions

1. Define death.
2. What are the stages experiences by dying person?
3. How do you confirm the occurrence of death?
4. What are the purposes of post mortem care?



GLOSSARY

Abrasion	a scraping or rubbing off of the skin.
Airborne precaution	precaution taken when a person has an illness that can be carried in the air or in the dust particles. Common measures include special air handling and ventilation.
Ambulatory	Walking
Ampule	small, glass sealed flask, often containing medication.
Anesthesia	complete or partial loss of sensation.
Anuria	complete suppression of urine secretion in the kidney.
Apex	lower point of the heart, formed by the tip of the left ventricle
Apical pulse	pulse normally heard at the heart's apex, which usually give the most accurate assessment of pulse rate
Aspiration	Inhalation of foodstuff, vomitus or saliva into the lungs.
Axilla	Armpit (under arm).
Autoclave	Equipment that decontaminates materials by exposing them to steam under pressure.
Apnea	Absence or lack of breathing
Anoxia	Lack of oxygen in the tissue.
Asphyxia	A condition produced by prolonged lack of oxygen

Asepsis	Absolute freedom from all microorganisms
Antiseptic	Harmless chemicals that can kill microorganisms or prevent them from multiplying.
Aplastic anemia	Anemia resulting from destruction of bone marrow cells.
Atelectasis	collapse of all or part of the lung.
Aseptic technique	Procedure used to prevent microorganisms from reaching the operation site.
Auscultation	externally hearing sounds from within the body to differentiate abnormal conditions.
Autopsy	examination of the body after death.
Base of support	balance or stability provided by the feet and their position.
Bed cradles	a wire or wooden frame placed over the patient's body or feet to support the weight of the bedclothes.
Blood pressure	The force exerted by the heart to pump the blood around the body
Bradycardia	Abnormally slow heartbeat.
Bradypnea	abnormally slow breathing, below 10 per minute.
Brand name	copyright name assigned by a company that makes the medication; also called the trade name.
Brain death	irreversible cessation of brain and brain stem function to the extent that

	cardiopulmonary function must be mechanical maintained.
Bounding pulse	Stronger than normal heartbeat.
Body mechanics	use of safe and efficient methods of moving and lifting.
Carotid pulse	pulse felt on either sides of the neck, over the carotid artery.
Capsule	a small gelatinous case for holding a dose of medicine; a membranous structuring enclosing another body structure, as the articular capsule in a joint.
Catheter	A soft rubber tube which is used for passage of fluid.
Center of gravity	the center of one's weight; half of one's body weight is below and half above, and half to the left and half to the right of the center of gravity.
Chemical name	medication name that describes its chemical composition (often same as generic name).
Cheyne-Stokes respiration:	breathing characterized by deep

Cyanosis	Bluish color of lips, tip of the nose, and ear lobes due to lack of or shortage of oxygen in the blood.
Cast	A material that supported an injured part of the body and makes it immobilize.
Clips	Metallic materials that keep the skin together.
Closed bed	bed used when preparing a unit for a new client- an unoccupied bed.
Congestion	Hyperemia, accumulation of blood in a part of blood or fluid in a part of the body e.g., lung.
Contact precaution	precaution taken against disease that can be transmitted through direct contact between a susceptible host's body surface and an infected or colonized person.
Cystitis	inflammation of urinary bladder.
Dangling	positioning of a client so that he or she is sitting on the edge of the bed with legs down and feet supported by a footstool or the floor.
Debridement	removal of foreign, dead, and contaminated material from a wound, so as to expose healthy underlying tissue.
Decontamination	The process of rendering an item free from infection.
Defecation	Act of excreting feces from the rectum.

Detergent	A substance usually dissolved in water used as an aid for cleaning purposes.
Diagnosis	The decision regarding the nature of an illness, arrived at by clinical assessment of the patient and result of investigation.
Diarrhea	abnormal frequency and fluidity of discharge from the bowels.
Diastole	The resting phase of the heart during which it fills with blood.
Digitalis	A drug given to slow and strengthen the heartbeat.
Disinfectant	A chemical used to kill microorganisms.
Dorsal lithotomy	examination position in which the client is lying on his or her back with the feet in stirrups.
Dosage	an amount in a prescription that contains the dose and the scheduled time.
Dry heat	Air heated to high temperature by electricity and used for sterilizing purposes.
Droplet precaution	precautions taken to prevent the spread of diseases transmitted by microorganisms propelled through the air from an infected person and deposited on the host's eyes, nose or mouth.
Dyspnea	Difficulty in breathing.
Dysuria	difficult or painful urination or voiding.



Flatus	Gas in the intestines.
Footboard	A board placed at the foot of the bed to support the feet.
Footdrop	contracture deformity that prevents the client from putting the heel on the floor; results from improper positioning or anterior leg muscle paralysis. On his or her back with the head elevated.
Fowler's position	a position in which the client is lying.
Gait	manner or style of walking.
Gastrostomy	Making an artificial opening into the stomach through which the patient is fed by pouring nourishment through a tube directly into the stomach.
Generic name	name assigned by a drug's first manufacturer (often the chemical name).
Halitosis	bad breath.
Hemoglobin	the oxygen carrying pigment in blood that gives blood its red color.
Hypertension	High blood pressure.
Hypo tension	Low blood pressure.
Hypothermia	low body temperature.



Melena	passage of dark colored stools containing partially or fully digested blood, also used to mean abnormal blood in the stool or vomitus.
Micturation	passage of urine from the urinary bladder; also called voiding , urinating.
Mitered corner	A triangular fold made in bedclothes to hold them in place at the corners.
Necrosis	Death of tissue.
Nocturia	excessive voiding (urination) during the night.
Nits	The eggs of a louse.
Occupied bed	bed holding a client that is unable to get up as a result of his or her condition or generalized weakness.
Occult	hidden.
Oliguria	deficient urinary secretion or infrequent urination.
Ophthalmic	medications that are instilled or administration directly to the eye.
Oral	of or pertaining to the mouth
Orthopenia	difficult breathing relieved by seating or standing erect
Output	All fluid lost from the body.
Otic	of the ear.

Postoperative bed	bed prepared for a client who is returning from surgery or another procedure that requires transfer into the bed from a stretcher or wheelchair.
Prescription	written formula for preparing and administering medication.
Preoperative	Before an operation
Projectile vomiting	emesis expelled with great force.
Prone	positioning a client so that he or she is lying on the stomach.
Pressure ulcer	ulcerated sore often caused by prolonged pressure on a bony prominence or other area, especially if the client is allowed to lie in one position for an extended period. Also called decubitus ulcer. (formerly "called bedsore").
Protective device	piece of equipment, most often a vest or a belt, used to ensure the safety of the client (ie, helping client to remain in a chair without falling); also called a client reminder device.
Protective isolation	attempts to prevent harmful microorganisms from coming into contact with the client; also called reverse or neutropenic isolation
Pulse	The beat of the heart felt in the arteries.
Pulse deficit	the difference between apical pulse and radial pulse.

Pulse pressure	difference between systolic & diastolic pressure.
Puncture	a whole made by a pointed object; penetration.
Recumbent	lying down
Rotation	process of turning about an axis, as rotation of the hand of the fetus in preparation of delivery.
Respiration	Breathing rate.
Radial pulse	pulse measured above the radial artery on the inside of the wrist.
Rectal	of the rectum
Retention enema	An injection of fluid that is retained in the rectum for absorption into the blood stream.
Restraints	Devices that limit the patient's ability to move in order to protect him/her from injury.
Septic wound	Infection wound; a wound containing infective microorganisms.
Sitzbath	A warm soaking of the rectum and perineal area.
Splint	A device for immobilizing part of the body
Spore	The seeds of microorganisms, which are resistant to drying, heat, and disinfectants
Standard precaution	precautions designed for the care of all clients regardless of diagnosis or infection status.
Sterile	Specially treated so that all microorganisms are destroyed

Stethoscope	Instrument for magnifying sound
Specimen	A small amount of body excretion or body fluid that is sent to a laboratory for examination.
Sphygmomanometer	Blood pressure apparatus.
Suppository	Rectally administered cones containing a medication in the base that is soluble at body temperature.
Sutures	Materials that keep broken skin together.
Systole	Blood pressure period during the beating phase of the heartbeat during which blood is expelled from heart.
Sepsis	Presence of microorganisms.
Synergism	joint action of agents in which the combined effects is greater than the sum of the individual parts.
Tablet	a compressed, spherical forms of medication.
Temperature	Degree of heat.
Tachycardia	Abnormally fast heartbeat.
Tachypnea	conditions in which breaths are abnormally rapid, more than 20 per minute
Thermometer	An instrument used to measure temperature.
Topical	medication that are applied directly to the skin or mucus membranes.
Traction	exertion of a pulling force ; an apparatus attached to the client to maintain stability of




REFERENCES

Caroline, B., R., and Mary T. K, (2003). *Textbook of Basic Nursing*, Eighth Edition, Lippincott comp.

Craven,R.F.,& Hirnle, C.J. (2000). *Fundamentals of nursing:Human health and function* (3rd ed.). Philadelphia: Lippincott Williams & Wilkins.

Evelyn P. (1989). *A Gener3/5.582lq A 200MenNuenGen*.



Benjamin/Cummings Publishing Comp. Philadelphia.

Lynda J., Carpento.(1995). *Nursing Care Plan and Documentation*, (2nd ed.) J.B. Lippincott Comp. Philadelphia.

Margaret C.(1983). *Practical Nursing*, (13th ed.), Baillier Tindall, London

Mary E. Scholes. (1986). *Hand Book of Nursing Procedures*, Blackwell Scientific Publications, London.

Robyn R. (1995). *Manual of Home Health Nursing Procedures*, Mosby Comp. Boston.

Sandra F. S., Donna J. D., and Barbara C. M. (2004). *Clinical Nursing Skill. Basic to advanced skills*, (6th ed.) PEARSON,Prentice Hall, New Jersey.

Sorensen and Luck Mann's. (1994). *Basic Nursing, A psychological approach*, (3rd ed.) W.B. Saunders comp. Philadelphia.

Smith & Dugan,(1996). *Clinical Nursing Skills*, basic to advanced skills, (4th ed.), Appleton & Lange comp

Suzanne C. S., and Brenda G. B. (1995). Bruner and Suddarth's Textbook of *MEDICAL-SURGICAL NURSING*, (8th ed.), Lippincott comp

Swearingen & Howard,(1996). *Photo Atlas of Nursing Procedures*, (3rd ed.), Benjamin Comp. California.

W. Kereta (1998). *a patient care procedure manual* for mid level health workers. Ethiopia.

