THE

SUMMARY PROCEEDINGS

3rd ANNUAL MALARIA CONTROL PROGRAM REVIEW
Ethiopia and Nigeria

Held on February 24, 2012 at
The Carter Center, Atlanta, Georgia

September 2012



In 2011, The Carter Center surpassed the milestone of helping the Nigeria and Ethiopia Ministries of Health to distribute 10 million insecticide-treated bed nets to fight malaria and lymphatic filariasis since 2004. On this pin, the hatched line pattern and the color white, which respectively symbolize "heroic efforts" and "peace" in Nigeria, represent the millions of people now able to sleep peacefully, undisturbed by biting mosquitoes that transmit these diseases.

Design by Sherri Richards

TABLE OF CONTENTS

ACKNOWLEDGMENTS ii
ACRONYMS iii

EXECUTIVE SUMMARY 1-7
RECOMMENDATIONS 6

NIGERIA PRESENTATIONS 8-19
Nigeria National Malaria M.. 6

ACKNOWLEDGEMENTS

The Carter Center's Malaria Control Program extends its sincere gratitude for the support provided by partners and donors, including Chevron Corporation, Clarke Mosquito Control, the Bill & Melinda Gates Foundation, Sanofi-aventis, the Kingdom of Saudi Arabia, Starr

Malaria elimination

The MCP is committed to demonstrating the potential to interrupt malaria transmission in even the most challenging situations when current (imperfect) tools are implemented at scale. Rather than waiting for the development of new technologies to address this goal, we are working to find new ways to use existing technologies in a more aggressive manner and with a focus on stopping transmission altogether, rather than only reducing morbidity and mortality.

During this year's review meeting, discussions were framed by presentations that focused on the current global interest in malaria elimination, highlighting the challenges and potential strategies for achieving malaria elimination and establishing the context for the Center's malaria activities. Dr. Larry Slutsker (CDC) highlighted some of the key recommendations outlined in the malaria Eradication Research Agenda (malERA). His talk emphasized the need for improved surveillance, diagnostic tools, and strategies when shifting from malaria control (identifying and treating cases) to elimination (identifying and treating infections including asymptomatic, latent and imported ones) (See Annex 3). Dr. Rick Steketee (PATH/MACEPA) gave an overview of

states, bringing the total number of nets distributed in Nigeria with TCC-assistance to nearly 4.3 million. Mr. Sallau described advocacy by General Gowon to accelerate the distribution of 6.5 million nets in the four TCC-assisted states that had not yet begun mass campaigns. To encourage the consistent and appropriate use and care of bed nets where LLINs have been distributed, TCC Nigeria assisted Ebonyi state's Roll Back Malaria (RBM) team with the design and implementation of a community-based behavior change communications (BCC) strategy. At baseline, 74.3% of household owned at least one net, 83.8% of nets were hanging, 83.0% of nets were used the previous night, and 65.6% of people reported sleeping under a net the previous night. After six months of intervention, 100% of households owned at least one net, 95% of LLINs were observed to be hanging, 98% of LLINs were used the previous night, and 97% of people reported sleeping under a net the previous night– all significant improvements from baseline.

Mr. Sallau also presented results from surveys conducted in Plateau and Abia states which showed the highest malaria prevalence among children 5-14 years of age, a group not traditionally targeted by malaria interventions. Net ownership prior to the mass campaigns was 7.2% in Abia and 35.1% in Plateau. A separate coverage survey conducted by the Center in Plateau s

million cases of malaria (both confirmed and clinically diagnosed) were reported in Ethiopia, down from approximately 4 million cases in 2010.

Mr. Solomon Kibret summarized TCC-supported malaria control activities in Ethiopia. The MCP in Ethiopia provided support to the FMOH and the Regional Health Bureaus for malaria control in Amhara, Oromia, Southern Nations Nationalities and People's Region (SNNPR), Gambella, and Beneshangul-Gumuz regions. This support consisted of routine monitoring of LLIN ownership and use, supportive supervision to improve the quality of case management and epidemic monitoring at a total of 662 health facilities, training of health workers on malaria diagnostics and treatment and, in Amhara region, mass screening and treatment of fever cases in the context of MDA campaigns for trachoma (MalTra weeks). Including training for MalTra week campaigns, an estimated 7,801 health workers received malaria training in 2011. A total of 48,715 cases of malaria were treated through MalTra weeks VI (East Amhara, May 2011) and VII (West Amhara, November 2011). These campaigns reach an estimated 98% of Amhara's population annually.

In addition, TCC provided support for Ethiopia's 2011 Malaria Indicator Survey (MIS)—a standardized nationwide survey to measure: access to, coverage and use of key malaria control interventions; prevalence of malaria-related fever, parasitemia and anemia; and malaria knowledge, attitudes and practices of women of reproductive age. TCC staff from headquarters and from the Ethiopia office assisted with survey design and training of survey teams (more than 300 personnel), provided financial, logistical (vehicles), and supervisory support for the Amhara phase of the survey, and continue to provide technical assistance for summary, analysis, and writing. The survey targeted an estimated 11,025 households, making it the largest MIS survey conducted in any country to date.

Dr. Gregory Noland presented a newly-developed malaria surveillance tool that was developed

malar( )]TJE /P <</MCID 5 >>h1(t)1(i)-2

RECOMMENDATIONS

GENERAL

1. Apply lessons learned from other Carter Center-assisted elimination programs to malaria elimination.

2. Lead the way in testing innovative strategies that are not part of the current global approach to malaria elimination. These potentially include starting with the most challenging places first and conducting mass drug administration for malaria.

NIGERIA

1. The state MOHs should support distribution of the remaining 6.5 million nets in Imo, Abia, Edo and Delta states, with TCC providing assistance according to its ability.

2. Introduce community-based behavior change communications (BCC), LLIN monitoring and LLIN replacement activities in Plateau state using CDDs and other community volunteers. These activities should employ a modified version of the strategy employed in Ebonyi state in 2010 that is less costly and more sustainable.

3. Develop and pilot strategy for strengthening malaria surveillance in Plateau and Enugu states.

4. Complete analysis of data from surveys conducted in Plateau, Abia, Imo and Ebonyi, and prepare manuscripts for publication.

5. Disseminate summary of the proceedings of the March 2012 Abuja conference widely to encourage collaboration between the malaria control and lymphatic filariasis elimination programs. (See Annex 5)

6. Increase efforts to share information about malaria activities in Nigeria with the global malaria community.

7. Launch surveys in Enugu as funds permit.

ETHIOPIA

1. Implement elimination strategy in selected woredas of Amhara Region, in collaboration with the Amhara Regional Health Bureau and PATH/MACEPA.

2. Develop systematic guidelines for responding to alerts provided by enhanced surveillance

Nigeria National Malaria Control Program Report

Presentation by Mrs. Chioma Ajoh, National Coordinator, National Malaria Control Program, Federal Ministry of Health

Background

Nigeria alone accounts for a quarter of the malaria burden in Africa. Malaria is endemic in Nigeria, with seasonal peaks during the rainy season. Almost 100% of the population is at risk for malaria infection and approximately 50% of the population will experience at least one episode each year. Malaria accounts for an estimated 66% of all health facility attendance and is responsible for 30% of deaths among children and 11% of maternal mortality in Nigeria. The social and economic burdens of malaria are also significant: malaria reduces the GDP of Nigeria by approximately 1% annually and is the leading cause of absenteeism.

The vision of the Nigerian National Malaria Control Program (NMCP) is a malaria-free Nigeria. The 2009-2013 NMCP Strategic Plan set ambitious targets of 80% of households owning at least two insecticide-treate66y 1nM Mnseowin-28.825 (h)2(e)3(e)a(h)-1e0n2an66c2(,)-h0.475(J)-1<34(nd)-3(p)-1m-4(er)(sn)-1(n)-1e

Nigerian National Malaria Control Strategy: Progress update

The Nigerian NMCP strategic plan focuses on four primary interventions: long-lasting insecticidal nets (LLINs), indoor residual spraying (IRS), effective case management, and malaria prevention in pregnancy. The plan



Figure 2. Status of national LLIN distribution campaigns as of February 2012.

While work remains to be done and a number of challenges must still be addressed, the successful LLIN campaigns conducted thus far reflect strong partnerships at all levels of the health system: the federal ministry and its partner NGOs, the state ministries, the local governments, communities, and individual volunteers. Nigerians have gone to great lengths to bring nets to even the most remote and hard to access communities.

In the states where mass campaigns have been completed, the NMCP plans to initiate routine distribution through immunization and ANC clinics, as well as child health days, in order to fill remaining gaps and maintain coverage rates. Additional components of the mixed method distribution system developed to achieve and sustain high coverage with LLINs include free or subsidized community-based "mop-up" campaigns and the sale of subsidized or at-cost LLINs through retailers, institutions, and local manufacturers.

Indoor Residual Spraying (IRS), Larviciding, and Environmental Management

The Nigerian target for IRS coverage is to spray 20% of households in the country (about 6 million) by 2013. IRS was implemented on a pilot basis in **Plateau**, Borno, Lagos, Adamawa, and Ogun states, and scaled up in one LGA in each of the seven states supported by the World Bank (Akwa Ibom, **Anambra**, Bauchi, Gombe, Jigawa, Kano, and Rivers). Lagos state has scaled-up IRS in a total of 8 LGAs. Select villages in Ondo state have also received IRS with support from the United Nations Development Project (UNDP). States supported by the Millennium Village project are scheduled to scale-up IRS in 2012, as is the President's Malaria Initiative-supported state of **Nasarawa**. A total of 63,000 households had been sprayed at the time of the program review meeting. In preparation for the gradual scale-up of IRS nationwide, trainings on malaria entomology have been conducted for the following people:

Edo and Delta are also targeted for distribution in 2012. The Center is working to ensure that 6,588,719 LLINs are distributed in these four remaining states before the end of 2012.

The continued delay of net distribution in these four states is currently the greatest challenge facing the Center's MCP in Nigeria. In order to address barriers to distribution, the Center's Nigerian staff worked with the state and local MOH in Abia and Imo states to revise the campaign strategy in order to reduce the amount of funding required. In addition, TCC organized advocacy visits by General Yakubu Gowon to both states. The former head of state has a long history of partnering with TCC for advocacy efforts, with a total of 322 field visits (beginning in 1998) in 23 states that were conducted in the context of the successful Guinea Worm Eradication Program. In December 2011, the General, accompanied by representatives of the Federal Ministry of Health, WHO, UNICEF and The Carter Center Nigeria, met with the governors of both Imo and Abia states, as well as with other representatives of the state and local governments. The purpose of the visits was to mobilize support and encourage the release of state funds for LLIN distribution. As the result of these visits, the state and local governments agreed to support the distribution of over 3 million LLINs to the two states in 2012.

Behavior Change Communications (BCC)

In 2011, TCC assisted the Ebonyi State Roll Back Malaria (RBM) program with the implementation of a new community-based behavior change communication strategy that was designed to increase rates of consistent and appropriate bed net use and care following the completion of the mass LLIN distribution campaign. The strategy, which was piloted in 6 sentinel villages of Ebonyi State, was informed by social behavioral theory and addressed a wide range of the possible determinants of net use.

Representatives of TCC and the state RBM team worked together to train a total of 87 volunteer community health promoters (CHPs) and 18 Nigerian supervisors in June 2011, and the project was launched in early July. The CHPs visited households on a monthly basis. During these visits, they assessed the household's current net ownership and maintenance behaviors, checking to see if the household owned enough nets, if the nets were hanging, if they were hanging at the appropriate height, and if they needed mending or washing. The CHPs also collected data on household net use. Subsequent interventions were tailored to fit the household's current behaviors. The CHPs engaged in conversations with household members to identify the specific barriers preventing them from adopting any recommended behaviors not yet practiced and then selected appropriate activities and messages to address those barriers. Specific targeted determinants of net use and appropriate care included access to nets, knowledge, attitudes, skills, self-efficacy, social support, structural barriers (e.g. not having a place to hang the net), and perceived social norms.

Figure 5. CHPs showing community members how to build portable frame for hanging bed nets over any sleeping space in Ebonyi State (photo by Adamu Sallau).

In addition to the monthly home visits, the CHPs organized community events, including community net mending and washing days, theatrical performances, net hanging

demonstrations, and workshops to show people how to use inexpensive, locally-available materials to build portable net-hanging frames. These frames can be used to hang a net over any sleeping space and can be easily moved out of the way during the day (Figure 5). The CHPs also organized meetings with community leaders to mobilize them in efforts to increase net use and appropriate net care.

The messages employed in the context of this project emphasized the effectiveness of bed nets for preventing the manifestations of lymphatic filariasis (LF) as well as malaria. Nigerian adults who consider malaria episodes as a minor illness (due to their having developed a degree of immunity to the infection) may have greater fear of developing "big legs" or enlarged testicles. Thus it is possible that messages linking bed nets to LF prevention will be more effective at encouraging universal net use by all members of the population, especially adult men.



Figure 6. Changes in LLIN hanging and use indicators after 6 months of intervention in six villages of Ebonyi state, Nigeria.

The net ownership data collected during monthly home visits motivated the Ebonyi state RBM program to conduct a "mop-up" campaign during which 1,094 LLINs were distributed to 554 households. By the end of the second month of the BCC intervention (August), 100% of the 1284 households in these six villages owned at least one LLIN (compared to 79.4% in July), and 93.4% owned at least two LLINs (up from 64.4% in July). After six months of intervention, there were significant improvements in all indicators related to net hanging and net use (Figure 6).

program's effects on intermediate outcomes such as malaria knowledge and attitudes, bed net

Net ownership was low in both states prior to the mass distribution campaigns, but was much higher in Plateau State where TCC

Ethiopia National Malaria Control Program

Presented by Ms. Hiwot Solomon, National Malaria Programme Focal Person, Federal Ministry of Health, Ethiopia

Malaria is a leading public health problem in Ethiopia. An estimated 55.7 million people (68% of the population) are at risk for malaria and around 80% of the 736 districts (woredas) in Ethiopia are considered "malarious." Malaria transmission is generally seasonal and unstable, though patterns and intensity of transmission vary throughout the country due to differences in altitude, rainfall and population movement. Protective immunity in Ethiopian populations is relatively low due to unstable transmission and, unlike large parts of sub-Saharan Africa, all age groups are at risk of infection and disease. P. falciparum

were printed and distributed to health posts. The objectives are that 100% of health facilities in epidemic prone areas adhere to the national epidemic and response plan; and 100% of health facilities and woreda(district)

Figure 11. Annual incidence of confirmed malaria (actual and predicted) and confirmed plus clinical malaria in Ethiopia, 1983-2011.

Health systems strengthening and capacity building. The aforementioned strategies for malaria control, prevention, and elimination in Ethiopia will be supported by additional cross-cutting measures that include: monitoring and evaluation, human resource development, financial management and

developed health worker capacity in epidemic monitoring and malaria case management. In addition, a total of 7,801 health workers were trained in RDT use and malaria case management in 2011.

Monitoring and evaluation. TCC conducts ongoing assessment of LLIN ownership, use, and care at the community level. Evaluation teams are comprised of the Carter Center's zone project coordinators, woreda health officers, and health extensions workers, all of whom are Ethiopian nationals. In 2011, a total of 7,018 households were assessed in Amhara, SNNPR, Oromia, Gambella, and Beneshangul-Gumuz regions of Ethiopia. As shown in Table 3, household net ownership was high (≥92%) in all surveyed regions except Oromia (74%). However, less than 50% of all household members slept under a net the previous night in all regions except Beneshangul-Gumuz (where the reported figure was 60%). There was considerable variation between regions in the proportion of households in which all nets

Enhanced System for Malaria Surveillance in Ethiopia

Presented by Dr. Gregory Noland, Malaria Epidemiologist, The Carter Center

Ethiopia has a strong history of routine surveillance systems for public health. Ethiopia has the further advantage of a high proportion of services being provided through the public sector. This is particularly important for Ethiopia where malaria is highly variable and characterized by an unstable transmission pattern in both space and time. In such an epidemiological context, routine health facility-based surveillance offers several advantages over cross-sectional surveys such as malaria indicator surveys (MIS) for collecting and analyzing malaria data because:

- It permits detailed information at community-level resolution
- It allows for monitoring of seasonal and interannual variations in indicators
- It allows tailored responses to case buildups or epidemics if there are no delays in reporting and analysis

However, issues such as under-reporting, lack of uniformity in reports, ambiguous interpretation of indicators, lack of information on reporting fraction (the % of health facilities reporting), and discrepancies between weekly reports and monthly summaries currently complicate the collection and analysis of malaria surveillance data in Ethiopia.

In order to address these needs, The Carter Center has developed an Excel-based "Enhanced Surveillance System" tool to record and monitor routinely collected MOH malaria data in Amhara Region—though the framework can be easily adapted to other locations. This platform was developed with unity in mind such that there is consistency in indicators and formatting across all TCC-assisted zones of Amhara. Importantly, it was also developed to allow harmonization with the Amhara Regional Health Bureau (RHB) malaria reporting systems, so that TCC zone project coordinators (ZPCs) can copy directly from data reported to FMOH Zonal Health Departments.

The system consists of a template file for recording weekly summary health facility data by district for each of the 10 zones within Amhara. In addition, a composite summary file is used to store and analyze data at for the entire region. An interactive "dashboard" page in the summary file allows the visualization of multiple malaria-related indicators on a single worksheet at district, zone, or regional level detail (Figure 13).

Following a 6-month pilot in two zones of Amhara (West Gojjam and North Gondar), the Enhanced Surveillance System was deployed to all 10 zones of Amhara in December 2011.
The advantages of this system include:

- A format that allows easy entry of existing FMOH data—avoiding the introduction of an additional reporting form to already over-burdened health workers.
- Automatic visual representation of data for multiple malaria indicators.
- Prompt access by zone-level ZPCs, who can immediately interpret and react to results.

We plan to continue developing and improving the system in partnership with TCC Ethiopia personnel and Amhara RHB colleagues; to potentially expand to other TCC-assisted malaria zones outside of Amhara; and also to adapt it to malaria surveillance in Nigeria in 2012.



Figure 13. Screen-shot of the enhanced surveillance system "dashboard" page

Annex 1: Overview of Malaria Disease, the History of The Carter Center Malaria Control Program, and Malaria Program Priorities

The Disease

Malaria is a parasitic disease caused by the single-celled organism *Plasmodium*, which infects the human liver and red blood cells. It is transmitted from person to person by the bite of the *Anopheles* mosquito,

wide scale-up of malaria control activities, with a focus on the mass distribution of long-lasting insecticidal nets (LLINs).

A dedicated Carter Center MCP was launched in Ethiopia in February 2006. The Ethiopian Minister of Health, Dr. Tedros Adhanom, requested that TCC join his country's national effort to provide protection to all 50 million Ethiopians at risk for malaria through an ambitious plan to distribute long lasting insecticidal nets in all malarious areas by the end of 2007. The Center was also asked to help in national efforts to monitor and evaluate the progress and effectiveness of the national control program. Since 2006, the program has built on TCC's existing programmatic networks in parts of Ethiopia while working

adoption of recommended behaviors. In Ethiopia, these messages have been disseminated using a number of different channels, including radio and television spots, films, posters, brochures/pamphlets, theater, t-shirts and caps, and mass education sessions. Malaria health education is provided during azithromycin[8] distribution for trachoma and mass testing and treatment of malaria cases (MalTra Weeks). Malaria messages are also highlighted during the distribution of ivermectin for onchocerciasis control. In Nigeria, health education for malaria has been provided in association with the distribution of insecticide treated bed nets, as well as in the context of an intensive



Figure 14. Elimination framework.

Other factors aside from transmission levels to be considered include:

- Logistical feasibility
- Availability of woreda-level surveillance data
-

Annex 5: Editor's Report - National Conference on the Collaboration between the Lymphatic Filariasis Elimination and Malaria Control Programs, March 27-28, 2012

In March 2012, the Nigerian Federal Ministry of Health and The Carter Center co-hosted the first national conference to stop transmission of lymphatic filariasis (LF) and malaria in Nigeria. The meeting was held on March 27-28 at the Shehu Musa Yar'Adua Center in Abuja. The conference was chaired by General (Dr.) Yakubu Gowon (former Head of State of the Federal Republic of Nigeria). The Honorable Minister of Health was represented by the Permanent Secretary of Health, Mrs. Fatima Bamidele. Participants at the conference included representatives from the Federal Ministry of Health, The Carter Center (TCC) Atlanta and Nigeria offices, the Bill & Melinda Gates Foundation (BMGF), the World Health Organization (WHO), the African Program for Onchocerciasis Control (APOC), the Department for International Development (DFID), SightSavers, Vision 2020, Mission to Save the Helpless (MITOSATH), Helen Keller International (HKI), as well as the Nigerian Neglected Tropical Diseases (NTDs) steering committee chairperson, the Nigerian Non-Governmental Development Organization (NGDO) coalition chairman, and the state program managers for LF & malaria. The Nigerian press was also well-represented at the meeting.

The schedule of meeting sessions is presented in Table 5 below. The plenary sessions were moderated by Dr. Lance Gordon, Professor Adenike Abiose, Professor O. O. Kale, Dr. Y. J. Jiya, and Professor O. Akogun.

Table 5: List of Presentations and Presenters

Presentation Title	Presenter	Institutional Affiliation
The Imperative for the Integration of LF Elimination and Malaria Control Programmes	Dr. Frank Richards	TCC Atlanta
Global Overview of Malaria	Dr. Amy Patterson	TCC Atlanta
Global Overview of LF	Mr. Darin Evans	TCC Atlanta
National Overview of Malaria	Dr. Joel Akilah	National Malaria Control Programme
National Overview of LF	Mr. E. Davies	National LF Elimination Programme
The Carter Center Plateau Experience	Dr. Abel Eigege	TCC Nigeria
Behavior Change Communication	Mr. Adamu Sallau	TCC Nigeria
LF and Malaria: Strategies and Opportunities for Synergies	Dr. Lance Gorden	BMGF
Integrated Vector Management	Dr. Amy Patterson	TCC Atlanta
Effective implementation of LF Elimination in Endemic Countries: Monitoring and Evaluation	Professor O. Akogun	Federal University of Technology, Yola
Sustainable Management Training Center: Integration Experience	Dr. Abel Eigege	TCC Nigeria

On the second day of the meeting, participants broke out into small groups to develop recommendations pertaining to integration of malaria and LF activities in the following areas: monitoring and evaluation, policy and funding, behavior change communications, treatment scale-up and delivery, vector control scale-up and delivery, and community and community health workers.

At the end of the meeting, the following resolutions and recommendations were made:

1. As a follow up from the malaria (ML)/lymphatic filariasis (LF) conference in Abuja, Federal

Annex 6: Publications and Abstracts since 2007

Publications

1) Graves PM, Ngondi JM, Hwang J, Getachew A, Gebre T, Mosher AW, Patterson AE, et al. (2011). Factors associated with mosquito net use by individuals in households owning nets in Ethiopia. Malaria Journal10:354.

2) Ngondi JM, Graves PM, Gebre T, Mosher AW, Shargie EB, Emerson PM, and Richards FO. (2011). Which nets are being used: factors associated with mosquito net use in Amhara, Oromia and Southern Nations, Nationalities and Peoples' Regions of Ethiopia. Malaria Journal10:92

3) Shargie EB, Ngondi J, Graves PM, Getachew A, Hwang J, Gebre T, Mosher AW, Ceccato P, Endeshaw T, Jima D, Tadesse Z, Tenaw E, Reithinger R, Emerson PM, Richards FO, and Ghebreyesus TA. (2010). Rapid increase in ownership and use of long-lasting insecticidal nets and decrease in prevalence of malaria in three regional states of Ethiopia, 2006-2007. Journal of Tropical Medicine.Volume 2010, Article ID 750978, 12 pages doi:10.1155/2010/750978.

4) Endeshaw T, Graves PM**,** Shargie EB, Gebre T, Ayele B, Yohannes G, Zerihun M, Genet A, Melak B, Kebede A, Jima D, Tadesse Z, Ngondi J, Mosher AW, Richards FO, and Emerson PM. (2010). Comparison of Parascreen Pan/Pf, Paracheck Pf and light microscopy for detection of malaria among febrile patients, Northwest Ethiopia. Trans Roy Soc Trop Med Hyg 104: 467-474.

Annex 7: List of Participants

Bill & Melinda Gates Foundation
Erin Shutes, M.P.H.

Centers for Disease Control
Dr. Leo F. Weakland
Dr. Stephen Blount
Dr. Julie Gutman
Dr. Mark Eberhard
Dr. Kimberly Lindblade
Dr. Laurence (Larry) Slutsker
Dr. Gail Stennies

Council of State and Territorial Epidemiology
Ms. LaKesha Robinson

Department of Public Health and Primary Care
Dr. Jeremiah Ngondi

Emory University
Dr Peter Brown
Dr. Sita Ranchod-Nilsson
Dr. Deb McFarland

Ethiopia
Mr. Ayeligne Mulualem Tuafie - Amhara Regional Health Bureau
Mr. Asefaw Getachew
Mrs. Hiwot Solomon Taffese - Ministry of Health
Mr Solomon Kibret Birhanie - The Carter Center
Dr. Zerihun Tadesse Gebreselassie - The Carter Center
Mr. Aseged Taye Zeleke - The Carter Center
Mr. Mulat Zerihun Lemu - The Carter Center

Annex 8: Malaria Program Review Meeting Agenda

February 24, 2012
8:00